



INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁶ : A01N 63/00, 63/04, 59/00, 43/84, 37/10, C11D 3/386, 3/48	A1	(11) International Publication Number: WO 97/42825 (43) International Publication Date: 20 November 1997 (20.11.97)
(21) International Application Number: PCT/DK97/00205 (22) International Filing Date: 6 May 1997 (06.05.97) (30) Priority Data: 0559/96 9 May 1996 (09.05.96) DK 0785/96 15 July 1996 (15.07.96) DK (71) Applicant (for all designated States except US): NOVO NORDISK A/S [DK/DK]; Novo Allé, DK-2880 Bagsværd (DK). (72) Inventor; and (75) Inventor/Applicant (for US only): JOHANSEN, Charlotte [DK/DK]; Novo Nordisk a/s, Novo Allé, DK-2880 Bagsværd (DK). (74) Common Representative: NOVO NORDISK A/S; Corporate Patents, Novo Allé, DK-2880 Bagsværd (DK).		(81) Designated States: AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CU, CZ, DE, DK, EE, ES, FI, GB, GE, GH, HU, IL, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, TJ, TM, TR, TT, UA, UG, US, UZ, VN, YU, ARIPO patent (GH, KE, LS, MW, SD, SZ, UG), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, ML, MR, NE, SN, TD, TG). Published <i>With international search report.</i>
(54) Title: ANTIMICROBIAL PEROXIDASE COMPOSITIONS (57) Abstract: <p>Enzymatic compositions comprising a <i>Coprinus</i> peroxidase, hydrogen peroxide or a source of hydrogen peroxide, and an enhancing agent such as an electron donor, e.g. phenothiazine-10-propionic acid; 2,2'-azino-bis(3-ethylbenzothiazoline-6-sulfonate); acetosyringate; C₁₋₈-alkylsyringate; or a water-soluble halide or thiocyanate salt such as potassium iodide, have antimicrobial properties useful e.g. for inhibiting or killing microorganisms present in laundry, on human or animal skin, hair, mucous membranes, oral cavities, teeth, wounds, bruises; and on hard surfaces; and can be used as disinfectant, a preservative for cosmetics, and for cleaning, disinfecting or inhibiting microbial growth on process equipment used for e.g. water treatment, food processing, chemical or pharmaceutical processing, paper pulp processing, and water sanitation.</p>		

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece	ML	Mali	TR	Turkey
BG	Bulgaria	HU	Hungary	MN	Mongolia	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MR	Mauritania	UA	Ukraine
BR	Brazil	IL	Israel	MW	Malawi	UG	Uganda
BY	Belarus	IS	Iceland	MX	Mexico	US	United States of America
CA	Canada	IT	Italy	NE	Niger	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NL	Netherlands	VN	Viet Nam
CG	Congo	KE	Kenya	NO	Norway	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NZ	New Zealand	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	PL	Poland		
CM	Cameroon	KR	Republic of Korea	PT	Portugal		
CN	China	KZ	Kazakhstan	RO	Romania		
CU	Cuba	LC	Saint Lucia	RU	Russian Federation		
CZ	Czech Republic	LI	Liechtenstein	SD	Sudan		
DE	Germany	LK	Sri Lanka	SE	Sweden		
DK	Denmark	LR	Liberia	SG	Singapore		
EE	Estonia						

ANTIMICROBIAL PEROXIDASE COMPOSITIONS

The present invention relates to an enzymatic composition
5 capable of killing or inhibiting microbial cells or
microorganisms, more specifically microbial cells or
microorganisms present in laundry, on hard surface, on
skin, teeth or mucous membranes; and for preserving food
products, cosmetics, paints, coatings, etc., the compo-
10 sition comprising a peroxidase enzyme and an enhancing
agent acting as electron donor.

BACKGROUND OF THE INVENTION

15

Various enzymatic antimicrobial compositions are known in
the art. For instance, WO 94/04127 discloses stabilized
dentifrice compositions which are capable of producing
antimicrobially effective concentrations of hypothiocyanite
20 ions. The compositions contain an oxidoreductase capable of
producing hydrogen peroxide and a peroxidase enzyme capable
of oxidizing thiocyanate ions, which are normally present
in saliva, to antimicrobial hypothiocyanite ions. Suitable
peroxidases include lactoperoxidase, myeloperoxidase,
25 salivary peroxidase and chloroperoxidase.

In EP-A-0 500 387 enzymatic antimicrobial compositions are
disclosed comprising a haloperoxidase, e.g. myelo-
peroxidase, eosinophil oxidase, lactoperoxidase and
30 chloroperoxidase, which selectively binds to and inhibits
the growth of target microorganisms in the presence of
peroxide and halide.

WO 95/27046 discloses an antimicrobial composition
35 comprising a Vanadium chloroperoxidase, halide ions, and
hydrogen peroxide or a hydrogen peroxide-generating agent.

The object of the invention is to provide a composition for killing or inhibiting microbial cells, i.e. for disinfection or preservation, which is easy to use and an effective alternative to the known disinfecting and
5 preserving compositions and methods.

SUMMARY OF THE INVENTION

10 Surprisingly, it has been found that the combined action of a peroxidase enzyme from the fungus *Coprinus* and an enhancing agent acting as electron-donor, when applied to e.g. a hard surface, skin, mucous membranes, oral cavity, hair, or laundry in the presence of hydrogen peroxide,
15 results in a hitherto unknown synergistic antimicrobial effect.

Thus, based on these findings the present invention provides, in a first aspect, an enzymatic antimicrobial
20 composition comprising or consisting essentially of a peroxidase obtainable from or produced by the fungus *Coprinus*, an enhancing agent, and hydrogen peroxide or a source of hydrogen peroxide.

25 The composition of the invention is useful as antimicrobial ingredient wherever such an ingredient is needed, for example for the preservation of food, beverages, cosmetics, deodorants, contact lens products, food ingredients or enzyme compositions; as a disinfectant for use e.g. on human
30 or animal skin, hair, oral cavity, mucous membranes, wounds, bruises or in the eye; for killing microbial cells in laundry; and for incorporation in cleaning compositions or disinfectants for hard surface cleaning or disinfection.

35 Accordingly, in further aspects, the present invention provides a method of inhibiting microorganisms present in laundry, wherein the laundry is treated with a soaking, washing or rinsing liquor comprising this composition; a

method of inhibiting microbial growth on a hard surface,
wherein the surface is contacted with this composition; and
a method of killing microbial cells present on human or
animal skin, mucous membranes, teeth, wounds, bruises or in
5 the eye or inhibiting the growth thereof, wherein the cells
to be killed or inhibited or the skin, mucous membrane,
teeth, wound or bruise are/is contacted with this composi-
tion.

10

DETAILED DESCRIPTION OF THE INVENTION

The term "microbial cells" denotes bacterial or fungal
cells, and the term "microorganism" denotes a fungus, a
15 bacterium or a yeast.

The term "hard surface" as used herein relates to any
surface which is essentially non-permeable for microor-
ganisms. Examples of hard surfaces are surfaces made from
20 metal, e.g. stainless steel, plastics, rubber, board,
glass, wood, paper, textile, concrete, rock, marble, gypsum
and ceramic materials which optionally may be coated, e.g.
with paint, enamel and the like. The hard surface can also
be a process equipment member of a cooling tower, a water
25 treatment plant, a dairy, a food processing plant, a
chemical or pharmaceutical process plant. Accordingly, the
composition according to the present invention is useful in
a conventional cleaning-in-place (C-I-P) system.

30 In the present context, the term "bactericidal" is to be
understood as capable of killing bacterial cells.

In the present context, the term "bacteriostatic" is to be
understood as capable of inhibiting bacterial growth, i.e.
35 inhibiting growing bacterial cells.

In the present context, the term "fungicidal" is to be un-
derstood as capable of killing fungal cells.

In the present context, the term "fungistatic" is to be understood as capable of inhibiting fungal growth, i.e. inhibiting growing fungal cells.


5

Without being bound to this theory, it is believed that the key reaction in the antimicrobial effect of the combined peroxidase/enhancing agent system of the present invention is the oxidation of essential protein and enzyme sulphydryl
10 groups.

The peroxidase enzyme is able to catalyse H_2O_2 -dependent oxidation of an electron-donor, e.g. halide ions or the thiocyanate ion (SCN^- , a pseudohalide) to yield halogens or
15 other oxidising agents. The oxidising agents make an electrophilic attack on microbial components, resulting in chemical modification of essential enzymes, transport systems, and other functional components. Sulphydryl groups are especially susceptible to electrophilic attack, and are
20 usually present in higher amounts than other easily oxidised groups. Aromatic amino acid residues are also susceptible to attack. Most aspects of antimicrobial action can be correlated with chemical modification of these nucleophilic components. Antimicrobial activity is favoured
25 by influences that increase the stability of the oxidising agent, provided that these influences do not interfere with their electrophilic character, or their ability to penetrate microbial membranes. Although H_2O_2 itself is a powerful oxidising agent, the H_2O_2 molecule is stabilised
30 and reacts slowly with biological materials. Also, most cells have enzymes that rapidly eliminate H_2O_2 . Peroxidase-catalysed oxidation of e.g. halides or SCN^- conserves the oxidising power of H_2O_2 in forms that react more rapidly, and for which the target cells may have no defense.

35 (Thomas, E.L. in "The Lactoperoxidase System". Ed. By Pruitt, K.M., and Tenovuo, J.O., New York, 1985).

The reaction catalysed by peroxidase can be written as



5 where AH_2 and A are reduced and oxidised forms of suitable electron donors; or, in case of halides or thiocyanate,



10

The hydrogen peroxide may be generated by an oxidoreductase enzyme and a substrate specific to that enzyme, in particular by those oxidoreductases which utilise water as a co-reactant and oxygen as an electron donor. Suitable
15 oxidoreductases include glucose oxidase, galactose oxidase, glycollate oxidase, lactate oxidase, L-gulonolactone oxidase, L-2-hydroxyacid oxidase, aldehyde oxidase, xanthine oxidase, D-aspartate oxidase, L-amino acid oxidase, D-amino acid oxidase, monoamine oxidase,
20 pyridoxaminephosphate oxidase, diamine oxidase, and sulfite oxidase. Glucose oxidase is most preferred. Suitable substrates are specific to the particular oxidoreductases chosen and are well known to the skilled person. For example, beta-D-glucose is a specific substrate for glucose
25 oxidase. Other suitable substrates include, but are not limited to D-glucose, D-galactose, L-sorbose, ethanol, tyramine, 1,4-diaminobutane, 2-aminophenol, glycollate, L-lactate, 2-deoxy-D-glucose, L-gulonolactone, L-galaconolactone, D-mannonolactone, L-2-hydroxyisocaproate,
30 acetaldehyde, butyraldehyde, xanthine, D-aspartate, D-glutamate, L-amino acids and D-amino acids.

It may be advantageous to use enzymatically generated hydrogen peroxide, since this source results in a
35 relatively low concentration of hydrogen peroxide under the biologically relevant conditions. Low concentrations of hydrogen peroxide result in an increase in the rate of

peroxidase-catalysed reaction.

The hydrogen peroxide can also be added to the composition per se or can be generated by perborate or percarbonate salts, preferably sodium percarbonate or sodium perborate.

The enzyme

- 10 The peroxidase employed in the method of the invention is preferably producible by plants (e.g. horseradish or soybean peroxidase) or microorganisms such as fungi or bacteria, more preferably by fungi including strains belonging to the subdivision Basidiomycotina, class
- 15 Basidiomycetes, especially the genus *Coprinus*, in particular *Coprinus cinereus* f. *microsporus* (IFO 8371), or *Coprinus macrorhizus*.

The peroxidase enzyme to be used in the method of the invention may be a monocomponent (recombinant) enzyme, i.e. enzymes essentially free from other proteins or enzyme proteins. A recombinant enzyme may be cloned and expressed according to standard techniques conventional to the skilled person. However, the enzyme may also be used in the

25 form of an enzyme preparation optionally enriched in an enzyme exhibiting the desired enzyme activity as the major enzymatic component, e.g. a mono-component enzyme preparation.

- 30 Particularly, a recombinantly produced peroxidase is a peroxidase derived from a *Coprinus* sp., in particular *C. macrorhizus* or *C. cinereus* according to WO 92/16634, or a variant thereof, e.g., a variant as described in WO 94/12621. Accordingly, a useful recombinant peroxidase may
- 35 be produced by using a DNA construct comprising the DNA sequence shown in SEQ ID No. 1 encoding a *Coprinus* sp. peroxidase, or a suitable modification thereof.

Examples of suitable modifications of the DNA sequence are nucleotide substitutions which do not give rise to another amino acid sequence of the peroxidase, but which correspond to the codon usage of the host organism, into which the DNA construct is introduced or nucleotide substitutions which do give rise to a different amino acid sequence and therefore, possibly, a different protein structure which might give rise to a peroxidase mutant with different properties than the native enzyme. Other examples of possible modifications are insertion of one or more nucleotides into the sequence, addition of one or more nucleotides at either end of the sequence, or deletion of one or more nucleotides at either end or within the sequence.

The DNA construct encoding the peroxidase may be prepared synthetically by established standard methods, e.g. the phosphoamidite method described by S.L. Beaucage and M.H. Caruthers, Tetrahedron Letters 22, 1981, pp. 1859-1869, or the method described by Matthes et al., EMBO Journal 3, 1984, pp. 801-805. According to the phosphoamidite method, oligonucleotides are synthesized, e.g. in an automatic DNA synthesizer, purified, annealed, ligated and cloned in suitable vectors.

The DNA construct may also be of genomic or cDNA origin, for instance obtained by preparing a genomic or cDNA library and screening for DNA sequences coding for all or part of the peroxidase by hybridization using synthetic oligonucleotide probes in accordance with standard techniques (cf. Sambrook et al., Molecular Cloning: A Laboratory Manual, 2nd Ed., Cold Spring Harbor, 1989). In this case, a genomic or cDNA sequence encoding the peroxidase may be modified at a site corresponding to the site(s) at which it is desired to introduce amino acid substitutions, e.g. by site-directed mutagenesis using synthetic oligonucleotides encoding the desired amino acid sequence

for homologous recombination in accordance with well-known procedures.

Finally, the DNA construct may be of mixed synthetic and
5 genomic, mixed synthetic and cDNA or mixed genomic and cDNA
origin prepared by ligating fragments of synthetic, genomic
or cDNA origin (as appropriate), the fragments
corresponding to various parts of the entire DNA construct,
in accordance with standard techniques. The DNA construct
10 may also be prepared by polymerase chain reaction using
specific primers, for instance as described in US 4,683,202
or R.K. Saiki et al., Science 239, 1988, pp. 487-491.

The DNA construct is normally inserted into a recombinant
15 expression vector. This may be any vector which may
conveniently be subjected to recombinant DNA procedures,
and the choice of vector will often depend on the host
cell, into which it is to be introduced. Thus, the vector
may be an autonomously replicating vector, i.e. a vector
20 which exists as an extrachromosomal entity, the replication
of which is independent of chromosomal replication, e.g. a
plasmid. Alternatively, the vector may be one which, when
introduced into a host cell, is integrated into the host
cell genome and replicated together with the chromosome(s),
25 into which it has been integrated.

In the vector, the DNA sequence encoding the peroxidase
should be operably connected to a suitable promoter and
terminator sequence. The promoter may be any DNA sequence
which shows transcriptional activity in the host cell of
30 choice and may be derived from genes encoding proteins
either homologous or heterologous to the host cell.

Examples of suitable promoters are those indicated above.
The procedures used to ligate the DNA sequences coding for
the peroxidase, the promoter and the terminator,
35 respectively, and to insert them into suitable vectors are
well known to persons skilled in the art (cf., for
instance, Sambrook et al., op.cit.).

A host cell is transformed with the expression vector. The host cell is a cell of a filamentous fungus, and is preferably a cell of an Aspergillus sp. as indicated above.

5

The medium used to culture the transformed host cells may be any conventional medium suitable for growing filamentous fungi. The transformants are usually stable and may be cultured in the absence of selection pressure. However, if
10 the transformants are found to be unstable, a selection marker introduced into the cells may be used for selection. If hemin or a heme-containing material (e.g. hemoglobin or red blood cells) is added to the medium, the yield of heme protein may be significantly increased.

15

The mature heme protein secreted from the host cells may conveniently be recovered from the culture medium by well-known procedures including separating the cells from the medium by centrifugation or filtration, and precipitating
20 proteinaceous components of the medium by means of a salt such as ammonium sulphate, followed by chromatographic procedures such as ion exchange chromatography, affinity chromatography, or the like.

25 In the context of this invention, peroxidase acting compounds comprise peroxidase active fragments derived from cytochromes, haemoglobin or peroxidase enzymes, and synthetic or semisynthetic derivatives thereof, e.g. iron porphins, iron porphyrins, and iron phthalocyanine and
30 derivatives thereof.

Determination of peroxidase activity: 1 peroxidase unit (POXU) is the amount of enzyme that catalyzes the conversion of 1 μmol hydrogen peroxide per minute at the following analytical conditions: 0.88 mM hydrogen peroxide,
35 1.67 mM 2,2'-azinobis(3-ethylbenzothiazoline-6-sulfonate), 0.1 M phosphate buffer, pH 7.0, incubated at 30°C, photome-

trically followed at 418 nm; molecular extinction coefficient $\epsilon = 3.6 \times 10^4 \text{ M}^{-1}\cdot\text{cm}^{-1}$.

The peroxidase enzyme may be present in the composition of
5 the invention corresponding to 0.01-100 POXU per ml of ready-to-use liquid, i.e. of washing solution, disinfecting liquid, preserving liquid, foot bath etc.

10 The enhancing agents

In a preferred embodiment of the invention, the enhancing agent capable of acting as an electron-donor is a source of ionic iodide which may be enzymatically converted to iodine
15 when contacted with peroxidase enzyme in an aqueous solution for a time and under conditions sufficient to permit the conversion.

Iodine (I_2) is widely used as a disinfectant, for many
20 types of situations, for example as skin cleansers, for wound disinfection, contact lens cleaning and water sanitation, to mention a few. In addition, iodine is also useful in catalysts, as an animal feed additive, in pharmaceuticals, and as polymer precursor additives.
25 Although the I_2 -based system of disinfection is extremely effective, several factors limit the scope of directly applying I_2 . In particular, the storage, transportation and handling of I_2 are extremely hazardous, due to the chemicals involved in production and also due to the
30 toxicity of I_2 itself even in moderate concentrations. Generally, I_2 is obtained from natural sources, such as brine, by processes that utilise strong inorganic acids, chlorine gas, and other hazardous chemicals. Iodophores have been developed as I_2 carriers to replace simple I_2
35 solutions for industrial and domestic disinfection. In addition, binary systems capable of generating I_2 from an I^- salt and a chemical oxidant are also available. Both these systems create the need for disposal of large,

potentially toxic amounts of by-products. Another alternative to both industrially producing I_2 on a large scale, and to applying I_2 as a disinfectant, has been found in the peroxidase-based generation of I_2 (US 4,282,324; US 4,617,190; US 4,588,586; US 4,937,072; US 5,055,287; US 5,227,161; US 5,169,455; US 4,996,146; US 4,576,817). Such methods involve the use of a peroxidase enzyme, the oxidising agent H_2O_2 , and a source of ionic iodide, Unfortunately, this method has the disadvantage of requiring the hazardous and volatile peroxide or peracid, which has to be either transported or generated *in situ* by additional enzymatic or chemical steps, this making the system more complex and/or costly.


In the present context, a preferred source of ionic iodide is a water-soluble iodide salt such as an alkaline metal iodide salt, e.g. potassium iodide (KI), sodium iodide (NaI), or lithium iodide, ammonium iodide, calcium iodide. Sodium iodide and potassium iodide are preferred.

20

Another preferred enhancing agent is a source of the thiocyanate ion (SCN^-), e.g. sodium thiocyanate, potassium thiocyanate, ammonium thiocyanate, and other thiocyanate salts, preferably sodium thiocyanate and potassium

25 thiocyanate.

In another preferred embodiment, a useful enhancing agent is the compound described by the following formula:



in which formula X represents (-O-) or (-S-), and the substituent groups R^1 - R^9 , which may be identical or different, independently represents any of the following radicals: hydrogen, halogen, hydroxy, formyl, carboxy, and esters and salts hereof, carbamoyl, sulfo, and esters and salts hereof, sulfamoyl, nitro, amino, phenyl, C_1 - C_{14} -alkyl, C_1 - C_5 -alkoxy, carbonyl- C_1 - C_5 -alkyl, aryl- C_1 - C_5 -alkyl; which carbamoyl, sulfamoyl, and amino groups may furthermore be unsubstituted or substituted once or twice with a substituent group R^{10} ; and which phenyl may furthermore be unsubstituted or substituted with one or more substituent groups R^{10} ; and which C_1 - C_{14} -alkyl, C_1 - C_5 -alkoxy, carbonyl- C_1 - C_5 -alkyl, and aryl- C_1 - C_5 -alkyl groups may be saturated or unsaturated, branched or unbranched, and may furthermore be unsubstituted or substituted with one or more substituent groups R^{10} ;


which substituent group R^{10} represents any of the following radicals: halogen, hydroxy, formyl, carboxy and esters and salts hereof, carbamoyl, sulfo and esters and salts hereof, sulfamoyl, nitro, amino, phenyl, aminoalkyl, piperidino, piperazinyl, pyrrolidin-1-yl, C_1 - C_5 -alkyl, C_1 - C_5 -alkoxy; which carbamoyl, sulfamoyl, and amino groups may furthermore be unsubstituted or substituted once or twice with hydroxy, C_1 - C_5 -alkyl, C_1 - C_5 -alkoxy; and which phenyl may furthermore be substituted with one or more of the following radicals: halogen, hydroxy, amino, formyl, carboxy and esters and salts hereof, carbamoyl, sulfo and esters and salts hereof, and sulfamoyl; and which C_1 - C_5 -alkyl, and C_1 - C_5 -alkoxy groups may furthermore be saturated or unsaturated, branched or unbranched, and may furthermore be substituted once or twice with any of the following radicals: halogen, hydroxy, amino, formyl, carboxy and esters and salts hereof, carbamoyl, sulfo and esters and salts hereof, and sulfamoyl;

or in which general formula two of the substituent groups R^1 - R^9 may together form a group -B-, in which B represents any of the following the groups: $(-CHR^{10}-N=N-)$, $(-CH=CH-)_n$, $(-CH=N-)_n$ or $(-N=CR^{10}-NR^{11}-)$, in which groups n-represents

an integer of from 1 to 3, R^{10} is a substituent group as defined above and R^{11} is defined as R^{10} . (It is to be understood that if the above mentioned formula comprises two or more R^{10} -substituent groups, these R^{10} -substituent groups may be the same or different).

In particular embodiments, the enhancing agent is 10-methylphenothiazine, phenothiazine-10-propionic acid, N-hydroxysuccinimide phenothiazine-10-propionate, 10-ethyl-phenothiazine-4-carboxylic acid, 10-ethylphenothiazine, 10-propylphenothiazine, 10-isopropylphenothiazine, methylphenothiazine-10-propionate, 10-phenylphenothiazine, 10-allylphenothiazine, 10-(3-(4-methylpiperazin-1-yl)propyl)-phenothiazine, 10-(2-pyrrolidin-1-yl-ethyl)phenothiazine, 2-methoxy-10-methyl-phenothiazine, 1-methoxy-10-methylphenothiazine, 3-methoxy-10-methylphenothiazine, 3,10-dimethylphenothiazine, 3,7,10-trimethylphenothiazine, 10-(2-hydroxyethyl)phenothiazine, 10-(3-hydroxypropyl)phenothiazine, 3-(2-hydroxyethyl)-10-methylphenothiazine, 3-hydroxymethyl-10-methylphenothiazine, 3,7-dibromophenothiazine-10-propionic acid, phenothiazine-10-propionamide, chlorpromazine, 2-chloro-10-methylphenothiazine, 2-acetyl-10-methylphenothiazine, 10-methylphenoxazine, 10-ethyl-phenoxazine, phenoxazine-10-propionic acid, 10-(2-hydroxyethyl)phenoxazine or 4-carboxyphenoxazine-10-propionic acid.

Another example of a useful enhancing agent is a compound described by the following formula:



in which formula A is a group such as -D, -CH=CH-D, -CH=CH-CH=CH-D, -CH=N-D, -N=N-D, or -N=CH-D, in which D is selected from the group consisting of -CO-E, -SO₂-E, -N-XY, and -N⁺-XYZ, in which E may be -H, -OH, -R, or -OR, and X and Y and Z may be identical or different and selected from -H and -R; R being a C₁-C₁₆ alkyl, preferably a C₁-C₈ alkyl, which alkyl may be saturated or unsaturated, branched or unbranched and optionally substituted with a carboxy, sulfo or amino group; and B and C may be the same or different and selected from C_mH_{2m+1}; 1 ≤ m ≤ 5.

In a preferred embodiment A in the above mentioned formula is -CO-E, in which E may be -H, -OH, -R, or -OR; R being a C₁-C₁₆ alkyl, preferably a C₁-C₈ alkyl, which alkyl may be saturated or unsaturated, branched or unbranched and optionally substituted with a carboxy, sulfo or amino group; and B and C may be the same or different and selected from C_mH_{2m+1}; 1 ≤ m ≤ 5.

In the above mentioned formula A may be placed meta to the hydroxy group instead of being placed in the paraposition as shown.


In particular embodiments, the enhancing agent is acetosyringone, methylsyringate, ethylsyringate, propylsyringate, butylsyringate, hexylsyringate, or octylsyringate.

Yet another useful enhancing agent is an azino compound described by the general formula




in which formula the symbols A and B, which may be identical or different, independently represent any of the substituents II, III, IV, and V,


(II)




(III)



(IV)



(V)



in which substituents the symbols X and Y, which may be identical or different, independently represent carbon, nitrogen, which nitrogen may be unsubstituted or substituted with a substituent group R^5 , sulfur, oxygen, selenium or tellurium;

and in which substituents the substituent groups R^1 , R^2 , R^3 , and R^4 , which may be identical or different, independently represent hydrogen, halogen, a hydroxy group, a C_1 - C_3 alkoxy group, a formyl group, a carboxy group, a sulfo group, a nitro group, a C_1 - C_5 alkyl group, which alkyl group may furthermore be saturated or unsaturated, linear or branched, or an amino group, which amino group may furthermore be unsubstituted or substituted once or twice with a substituent group R^5 ;

which substituent group R^5 represents halogen, a hydroxy group, a C_1 - C_3 alkoxy group, a C_1 - C_5 alkyl group, or an

amino group.

The peroxidase enhancing agent may be in free form or in the form of an addition salt.

5

In preferred embodiments, the substituent groups R^1 , R^2 , R^3 , and R^4 , which may be identical or different, independently represent hydrogen, halogen, a hydroxy group, a C_1 - C_3 alkyl group, or a sulfo group. Preferably, the halogen is fluoro, 10 chloro, or bromo. Preferably, the C_1 - C_3 alkyl group is methyl, ethyl, propyl, or isopropyl.

In preferred embodiments, the substituent group R^5 represents halogen, a hydroxy group, a C_1 - C_3 alkoxy group, a C_1 - 15 C_3 alkyl group, or an amino group.

In a most preferred embodiment, a peroxidase enhancing agent of the invention is 2,2'-azino-bis(3-ethyl-benzothiazoline-6-sulfonate). This compound, abbreviated 20 ABTS, is a chromogenic substrate, and a common peroxidase and phenol oxidase assay agent.

It has, moreover, been demonstrated that ABTS, contrary to the enhancers known and described above, is capable of 25 acting as a peroxidase enhancing agent at highly alkaline conditions, i.e. above pH 9. This feature allows ABTS to be implemented into e.g. detergent compositions, intended for performance in the range pH 7-13, particularly the range pH 8-12, preferably the range pH 9-11.

30

The enhancing agent may be present in the antimicrobial composition in concentrations corresponding to from 0.005 to 1000 μ mole per g of substrate (microbial cells, 35 biomass), preferably 0.05 to 500 μ mole per g of substrate, more preferably 0.5 to 100 μ mole per g of substrate.

Stability of the Radical of the Enhancing Agent

Without being limited to any theory it is presently contemplated that there is a positive correlation between
5 the half-life of the radical which the enhancing agent forms in the relevant aqueous medium and its efficiency, and that this half-life is significantly longer than the half-life of any of the substances selected from the group consisting of p-hydroxycinnamic acid, 2,4-dichlorophenol,
10 p-hydroxybenzene sulphonate, vanillin and p-hydroxybenzoic acid (i.e. the enhancing agents disclosed in WO 92/18683).

As the half-life of the radical is dependent on, inter alia, the pH, the temperature and the buffer of the aqueous
15 medium, it is very important that all these factors are the same when the half-lives of the radicals of various enhancing agents are compared.

The composition

20

The enzymatic composition of the invention may further comprise auxiliary agents such as wetting agents, thickening agents, buffer, stabilisers, perfume, colourants, fillers and the like.

25

Useful wetting agents are surfactants, i.e. non-ionic, anionic, amphoteric or zwitterionic surfactants. Examples of useful surfactants are mentioned below under "Uses".

30 The composition of the invention may be used in the form of a powder which is to be dissolved in water prior to use, or may be a gelled product or a liquid product. The composition may be a concentrated product or a ready-to-use product.

35

In use, the concentrated product is typically diluted with water to provide a medium having an effective antimicrobial activity, applied to the object to be disinfected or

preserved, and allowed to react with the microorganisms present.

The optimum pH condition is usually a compromise between optimum stability and optimum activity of the peroxidase enzyme, optimum stability and optimum reactivity (oxidation potential) of the radical of the enhancing agent, and the choice of buffering system.

10

Uses

The composition of the invention may be incorporated into a detergent or cleaning composition comprising more enzyme types useful in detergent or cleaning compositions, preferably at least one further enzyme selected from the group consisting of proteases, amylases, cutinases, peroxidases, oxidases, laccases, cellulases, xylanases, and lipases.

20

Surfactant system

The detergent compositions according to the present invention comprise a surfactant system, wherein the surfactant can be selected from nonionic and/or anionic and/or cationic and/or ampholytic and/or zwitterionic and/or semi-polar surfactants.

30 The surfactant is typically present at a level from 0.1% to 60% by weight.

The surfactant is preferably formulated to be compatible with enzyme components present in the composition. In liquid or gel compositions the surfactant is most preferably formulated in such a way that it promotes, or at least does not degrade, the stability of any enzyme in these compositions.

Preferred systems to be used according to the present invention comprise as a surfactant one or more of the nonionic and/or anionic surfactants described herein.

5

Polyethylene, polypropylene, and polybutylene oxide condensates of alkyl phenols are suitable for use as the nonionic surfactant of the surfactant systems of the present invention, with the polyethylene oxide condensates being preferred. These compounds include the condensation products of alkyl phenols having an alkyl group containing from about 6 to about 14 carbon atoms, preferably from about 8 to about 14 carbon atoms, in either a straight chain or branched-chain configuration with the alkylene oxide. In a preferred embodiment, the ethylene oxide is present in an amount equal to from about 2 to about 25 moles, more preferably from about 3 to about 15 moles, of ethylene oxide per mole of alkyl phenol. Commercially available nonionic surfactants of this type include IgepalTM CO-630, marketed by the GAF Corporation; and TritonTM X-45, X-114, X-100 and X-102, all marketed by the Rohm & Haas Company. These surfactants are commonly referred to as alkylphenol alkoxylates (e.g., alkyl phenol ethoxylates).

25 The condensation products of primary and secondary aliphatic alcohols with about 1 to about 25 moles of ethylene oxide are suitable for use as the nonionic surfactant of the nonionic surfactant systems of the present invention. The alkyl chain of the aliphatic alcohol can either be straight or branched, primary or secondary, and generally contains from about 8 to about 22 carbon atoms. Preferred are the condensation products of alcohols having an alkyl group containing from about 8 to about 20 carbon atoms, more preferably from about 10 to about 18 carbon atoms, with from about 2 to about 10 moles of ethylene oxide per mole of alcohol. About 2 to about 7 moles of ethylene oxide and most preferably from 2 to 5 moles of ethylene oxide per mole of alcohol are present in


said condensation products. Examples of commercially available nonionic surfactants of this type include TergitolTM 15-S-9 (The condensation product of C₁₁-C₁₅ linear alcohol with 9 moles ethylene oxide), TergitolTM 24-L-6 NMW 5 (the condensation product of C₁₂-C₁₄ primary alcohol with 6 moles ethylene oxide with a narrow molecular weight distribution), both marketed by Union Carbide Corporation; NeodolTM 45-9 (the condensation product of C₁₄-C₁₅ linear alcohol with 9 moles of ethylene oxide), NeodolTM 23-3 (the 10 condensation product of C₁₂-C₁₃ linear alcohol with 3.0 moles of ethylene oxide), NeodolTM 45-7 (the condensation product of C₁₄-C₁₅ linear alcohol with 7 moles of ethylene oxide), NeodolTM 45-5 (the condensation product of C₁₄-C₁₅ linear alcohol with 5 moles of ethylene oxide) marketed by 15 Shell Chemical Company, KyroTM EOB (the condensation product of C₁₃-C₁₅ alcohol with 9 moles ethylene oxide), marketed by The Procter & Gamble Company, and Genapol LA 050 (the condensation product of C₁₂-C₁₄ alcohol with 5 moles of ethylene oxide) marketed by Hoechst. Preferred 20 range of HLB in these products is from 8-11 and most preferred from 8-10.

Also useful as the nonionic surfactant of the surfactant systems of the present invention are alkylpolysaccharides 25 disclosed in US 4,565,647, having a hydrophobic group containing from about 6 to about 30 carbon atoms, preferably from about 10 to about 16 carbon atoms and a polysaccharide, e.g. a polyglycoside, hydrophilic group containing from about 1.3 to about 10, preferably from 30 about 1.3 to about 3, most preferably from about 1.3 to about 2.7 saccharide units. Any reducing saccharide containing 5 or 6 carbon atoms can be used, e.g., glucose, galactose and galactosyl moieties can be substituted for the glucosyl moieties (optionally the hydrophobic group is 35 attached at the 2-, 3-, 4-, etc. positions thus giving a glucose or galactose as opposed to a glucoside or galactoside). The intersaccharide bonds can be, e.g., between the one position of the additional saccharide units

and the 2-, 3-, 4-, and/or 6- positions on the preceding saccharide units.

The preferred alkylpolyglycosides have the formula

5



wherein R^2 is selected from the group consisting of alkyl, alkylphenyl, hydroxyalkyl, hydroxyalkylphenyl, and mixtures
10 thereof in which the alkyl groups contain from about 10 to about 18, preferably from about 12 to about 14, carbon atoms; n is 2 or 3, preferably 2; t is from 0 to about 10, preferably 0; and x is from about 1.3 to about 10, preferably from about 1.3 to about 3, most preferably from
15 about 1.3 to about 2.7. The glycosyl is preferably derived from glucose. To prepare these compounds, the alcohol or alkylpolyethoxy alcohol is formed first and then reacted with glucose, or a source of glucose, to form the glucoside (attachment at the 1-position). The additional glycosyl
20 units can then be attached between their 1-position and the preceding glycosyl units 2-, 3-, 4-, and/or 6-position, preferably predominantly the 2-position.

The condensation products of ethylene oxide with a
25 hydrophobic base formed by the condensation of propylene oxide with propylene glycol are also suitable for use as the additional nonionic surfactant systems of the present invention. The hydrophobic portion of these compounds will preferably have a molecular weight from about 1500 to about
30 1800 and will exhibit water insolubility. The addition of polyoxyethylene moieties to this hydrophobic portion tends to increase the water solubility of the molecule as a whole, and the liquid character of the product is retained up to the point where the polyoxyethylene content is about
35 50% of the total weight of the condensation product, which corresponds to condensation with up to about 40 moles of ethylene oxide. Examples of compounds of this type include certain of the commercially available PluronicTM

surfactants, marketed by BASF.

Also suitable for use as the nonionic surfactant of the nonionic surfactant system of the present invention, are the condensation products of ethylene oxide with the product resulting from the reaction of propylene oxide and ethylenediamine. The hydrophobic moiety of these products consists of the reaction product of ethylenediamine and excess propylene oxide, and generally has a molecular weight of from about 2500 to about 3000. This hydrophobic moiety is condensed with ethylene oxide to the extent that the condensation product contains from about 40% to about 80% by weight of polyoxyethylene and has a molecular weight of from about 5,000 to about 11,000. Examples of this type of nonionic surfactant include certain of the commercially available TetronicTM compounds, marketed by BASF.

Preferred for use as the nonionic surfactant of the surfactant systems of the present invention are polyethylene oxide condensates of alkyl phenols, condensation products of primary and secondary aliphatic alcohols with from about 1 to about 25 moles of ethyleneoxide, alkylpolysaccharides, and mixtures hereof. Most preferred are C₈-C₁₄ alkyl phenol ethoxylates having from 3 to 15 ethoxy groups and C₈-C₁₈ alcohol ethoxylates (preferably C₁₀ avg.) having from 2 to 10 ethoxy groups, and mixtures thereof.

Highly preferred nonionic surfactants are polyhydroxy fatty acid amide surfactants of the formula



wherein R¹ is H, or R¹ is C₁₋₄ hydrocarbyl, 2-hydroxyethyl, 2-hydroxypropyl or a mixture thereof, R² is C₅₋₃₁ hydrocarbyl, and Z is a polyhydroxyhydrocarbyl having a linear hydrocarbyl chain with at least 3 hydroxyls directly connected to the chain, or an alkoxylated derivative thereof. Preferably, R¹ is methyl, R² is straight C₁₁₋₁₅

alkyl or C₁₆₋₁₈ alkyl or alkenyl chain such as coconut alkyl or mixtures thereof, and Z is derived from a reducing sugar such as glucose, fructose, maltose or lactose, in a reductive amination reaction.


5

Highly preferred anionic surfactants include alkyl alkoxyated sulfate surfactants. Examples hereof are water soluble salts or acids of the formula RO(A)_mSO₃M wherein R is an unsubstituted C₁₀-C₂₄ alkyl or hydroxyalkyl group
10 having a C₁₀-C₂₄ alkyl component, preferably a C₁₂-C₂₀ alkyl or hydro-xyalkyl, more preferably C₁₂-C₁₈ alkyl or hydroxyalkyl, A is an ethoxy or propoxy unit, m is greater than zero, typically between about 0.5 and about 6, more preferably between about 0.5 and about 3, and M is H or a
15 cation which can be, for example, a metal cation (e.g., sodium, potassium, lithium, calcium, magnesium, etc.), ammonium or substituted-ammonium cation. Alkyl ethoxyated sulfates as well as alkyl propoxyated sulfates are contemplated herein. Specific examples of substituted
20 ammonium cations include methyl-, dimethyl, trimethyl-ammonium cations and quaternary ammonium cations such as tetramethyl-ammonium and dimethyl piperdinium cations and those derived from alkylamines such as ethylamine, diethylamine, triethylamine, mixtures thereof, and the
25 like. Exemplary surfactants are C₁₂-C₁₈ alkyl polyethoxylate (1.0) sulfate (C₁₂-C₁₈E(1.0)M), C₁₂-C₁₈ alkyl polyethoxylate (2.25) sulfate (C₁₂-C₁₈(2.25)M, and C₁₂-C₁₈ alkyl polyethoxylate (3.0) sulfate (C₁₂-C₁₈E(3.0)M), and C₁₂-C₁₈ alkyl polyethoxylate (4.0) sulfate (C₁₂-C₁₈E(4.0)M), wherein
30 M is conveniently selected from sodium and potassium. Suitable anionic surfactants to be used are alkyl ester sulfonate surfactants including linear esters of C₈-C₂₀ carboxylic acids (i.e., fatty acids) which are sulfonated with gaseous SO₃ according to "The Journal of the American
35 Oil Chemists Society", 52 (1975), pp. 323-329. Suitable starting materials would include natural fatty substances as derived from tallow, palm oil, etc.

The preferred alkyl ester sulfonate surfactant, especially for laundry applications, comprise alkyl ester sulfonate surfactants of the structural formula:

5

10



wherein R^3 is a C_8 - C_{20} hydrocarbyl, preferably an alkyl, or combination thereof, R^4 is a C_1 - C_6 hydrocarbyl, preferably an alkyl, or combination thereof, and M is a cation which forms a water soluble salt with the alkyl ester sulfonate. Suitable salt-forming cations include metals such as sodium, potassium, and lithium, and substituted or unsubstituted ammonium cations, such as monoethanolamine, diethanolamine, and triethanolamine. Preferably, R^3 is C_{10} - C_{16} alkyl, and R^4 is methyl, ethyl or isopropyl. Especially preferred are the methyl ester sulfonates wherein R^3 is C_{10} - C_{16} alkyl.

Other suitable anionic surfactants include the alkyl sulfate surfactants which are water soluble salts or acids of the formula $ROSO_3M$ wherein R preferably is a C_{10} - C_{24} hydrocarbyl, preferably an alkyl or hydroxyalkyl having a C_{10} - C_{20} alkyl component, more preferably a C_{12} - C_{18} alkyl or hydroxyalkyl, and M is H or a cation, e.g., an alkali metal cation (e.g. sodium, potassium, lithium), or ammonium or substituted ammonium (e.g. methyl-, dimethyl-, and trimethyl ammonium cations and quaternary ammonium cations such as tetramethyl-ammonium and dimethyl piperdinium cations and quaternary ammonium cations derived from alkylamines such as ethylamine, diethylamine, triethylamine, and mixtures thereof, and the like). Typically, alkyl chains of C_{12} - C_{16} are preferred for lower

wash temperatures (e.g. below about 50°C) and C₁₆-C₁₈ alkyl chains are preferred for higher wash temperatures (e.g. above about 50°C).

- 5 Other anionic surfactants useful for deterative purposes can also be included in the laundry detergent compositions of the present invention. These can include salts (including, for example, sodium, potassium, ammonium, and substituted ammonium salts such as mono- di- and triethanolamine salts)
- 10 of soap, C₈-C₂₂ primary or secondary alkanesulfonates, C₈-C₂₄ olefinsulfonates, sulfonated polycarboxylic acids prepared by sulfonation of the pyrolyzed product of alkaline earth metal citrates, e.g., as described in British patent specification No. 1,082,179, C₈-C₂₄ alkylpo-
- 15 lyglycolethersulfates (containing up to 10 moles of ethylene oxide); alkyl glycerol sulfonates, fatty acyl glycerol sulfonates, fatty oleyl glycerol sulfates, alkyl phenol ethylene oxide ether sulfates, paraffin sulfonates, alkyl phosphates, isethionates such as the acyl
- 20 isethionates, N-acyl taurates, alkyl succinamates and sulfosuccinates, monoesters of sulfosuccinates (especially saturated and unsaturated C₁₂-C₁₈ monoesters) and diesters of sulfosuccinates (especially saturated and unsaturated C₆-C₁₂ diesters), acyl sarcosinates, sulfates of
- 25 alkylpolysaccharides such as the sulfates of alkylpolyglucoside (the nonionic nonsulfated compounds being described below), branched primary alkyl sulfates, and alkyl polyethoxy carboxylates such as those of the formula $RO(CH_2CH_2O)_k-CH_2COO-M^+$ wherein R is a C₈-C₂₂ alkyl, k
- 30 is an integer from 1 to 10, and M is a soluble salt forming cation. Resin acids and hydrogenated resin acids are also suitable, such as rosin, hydrogenated rosin, and resin acids and hydrogenated resin acids present in or derived from tall oil.
- 35 Alkylbenzene sulfonates are highly preferred. Especially preferred are linear (straight-chain) alkyl benzene sulfonates (LAS) wherein the alkyl group preferably contains from 10 to 18 carbon atoms.

sulfonates (LAS) wherein the alkyl group preferably contains from 10 to 18 carbon atoms.

Further examples are described in "Surface Active Agents 5 and Detergents" (Vol. I and II by Schwartz, Perry and Berch). A variety of such surfactants are also generally disclosed in US 3,929,678, (Column 23, line 58 through Column 29, line 23, herein incorporated by reference).


10 When included therein, the laundry detergent compositions of the present invention typically comprise from about 1% to about 40%, preferably from about 3% to about 20% by weight of such anionic surfactants.

15 The laundry detergent compositions of the present invention may also contain cationic, ampholytic, zwitterionic, and semi-polar surfactants, as well as the nonionic and/or anionic surfactants other than those already described herein.

20

Cationic deterative surfactants suitable for use in the laundry detergent compositions of the present invention are those having one long-chain hydrocarbyl group. Examples of such cationic surfactants include the ammonium surfactants
25 such as alkyltrimethylammonium halogenides, and those surfactants having the formula:

30



35

wherein R^2 is an alkyl or alkyl benzyl group having from about 8 to about 18 carbon atoms in the alkyl chain, each R^3 is selected from the group consisting of $-\text{CH}_2\text{CH}_2-$, $-\text{CH}_2\text{CH}(\text{CH}_3)-$, $-\text{CH}_2\text{CH}(\text{CH}_2\text{OH})-$, $-\text{CH}_2\text{CH}_2\text{CH}_2-$, and mixtures thereof; each R^4 is selected from the group consisting of C_1 - C_4 alkyl, C_1 - C_4 hydroxyalkyl, benzyl ring structures formed by joining the two R^4 groups, $-\text{CH}_2\text{CHOHCHOHCO}R^6\text{CHOHCH}_2\text{OH}$, wherein R^6 is any hexose or hexose polymer having a molecular weight less than about 1000, and hydrogen when y is not 0; R^5 is the same as R^4 or is an alkyl chain, wherein the total number of carbon atoms or R^2 plus R^5 is not more than about 18; each y is from 0 to about 10, and the sum of the y values is from 0 to about 15; and X is any compatible anion.

15

Highly preferred cationic surfactants are the water soluble quaternary ammonium compounds useful in the present composition having the formula:

20




wherein R_1 is C_8 - C_{16} alkyl, each of R_2 , R_3 and R_4 is independently C_1 - C_4 alkyl, C_1 - C_4 hydroxy alkyl, benzyl, and $-(\text{C}_2\text{H}_4)_x\text{H}$ where x has a value from 2 to 5, and X is an anion. Not more than one of R_2 , R_3 or R_4 should be benzyl.

The preferred alkyl chain length for R_1 is C_{12} - C_{15} , particularly where the alkyl group is a mixture of chain lengths derived from coconut or palm kernel fat or is derived synthetically by olefin build up or OXO alcohols synthesis.

Preferred groups for R_2 , R_3 and R_4 are methyl and hydroxyethyl groups and the anion X may be selected from halide, methosulphate, acetate and phosphate ions.

Examples of suitable quaternary ammonium compounds of formulae (i) for use herein are:

- coconut trimethyl ammonium chloride or bromide;
 coconut methyl dihydroxyethyl ammonium chloride or
 bromide;
 decyl triethyl ammonium chloride;
 5 decyl dimethyl hydroxyethyl ammonium chloride or bromide;
 C₁₂₋₁₅ dimethyl hydroxyethyl ammonium chloride or bromide;
 coconut dimethyl hydroxyethyl ammonium chloride or
 bromide;
 myristyl trimethyl ammonium methyl sulphate;
 10 lauryl dimethyl benzyl ammonium chloride or bromide;
 lauryl dimethyl (ethenoxy)₄ ammonium chloride or bromide;
 choline esters (compounds of formula (i) wherein R₁ is



- di-alkyl imidazolines [compounds of formula (i)].
 20 Other cationic surfactants useful herein are also described
 in US 4,228,044 and in EP 000 224.

When included therein, the laundry detergent compositions
 25 of the present invention typically comprise from 0.2% to
 about 25%, preferably from about 1% to about 8% by weight
 of such cationic surfactants.

Ampholytic surfactants are also suitable for use in the
 30 laundry detergent compositions of the present invention.
 These surfactants can be broadly described as aliphatic
 derivatives of secondary or tertiary amines, or aliphatic
 derivatives of heterocyclic secondary and tertiary amines
 in which the aliphatic radical can be straight- or
 35 branched-chain. One of the aliphatic substituents contains
 at least about 8 carbon atoms, typically from about 8 to
 about 18 carbon atoms, and at least one contains an anionic
 water-solubilizing group, e.g. carboxy, sulfonate, sulfate.

See US 3,929,678 (column 19, lines 18-35) for examples of ampholytic surfactants.

When included therein, the laundry detergent compositions of the present invention typically comprise from 0.2% to about 15%, preferably from about 1% to about 10% by weight of such ampholytic surfactants.

Zwitterionic surfactants are also suitable for use in laundry detergent compositions. These surfactants can be broadly described as derivatives of secondary and tertiary amines, derivatives of heterocyclic secondary and tertiary amines, or derivatives of quaternary ammonium, quaternary phosphonium or tertiary sulfonium compounds. See US 3,929,678 (column 19, line 38 through column 22, line 48) for examples of zwitterionic surfactants.

When included therein, the laundry detergent compositions of the present invention typically comprise from 0.2% to about 15%, preferably from about 1% to about 10% by weight of such zwitterionic surfactants.

Semi-polar nonionic surfactants are a special category of nonionic surfactants which include water-soluble amine oxides containing one alkyl moiety of from about 10 to about 18 carbon atoms and 2 moieties selected from the group consisting of alkyl groups and hydroxyalkyl groups containing from about 1 to about 3 carbon atoms; watersoluble phosphine oxides containing one alkyl moiety of from about 10 to about 18 carbon atoms and 2 moieties selected from the group consisting of alkyl groups and hydroxyalkyl groups containing from about 1 to about 3 carbon atoms; and water-soluble sulfoxides containing one alkyl moiety from about 10 to about 18 carbon atoms and a moiety selected from the group consisting of alkyl and hydroxyalkyl moieties of from about 1 to about 3 carbon atoms.

Semi-polar nonionic detergent surfactants include the amine oxide surfactants having the formula:



wherein R^3 is an alkyl, hydroxyalkyl, or alkyl phenyl group or mixtures thereof containing from about 8 to about 22 carbon atoms; R^4 is an alkylene or hydroxyalkylene group containing from about 2 to about 3 carbon atoms or mixtures thereof; x is from 0 to about 3; and each R^5 is an alkyl or hydroxyalkyl group containing from about 1 to about 3 carbon atoms or a polyethylene oxide group containing from about 1 to about 3 ethylene oxide groups. The R^5 groups can be attached to each other, e.g., through an oxygen or nitrogen atom, to form a ring structure.

These amine oxide surfactants in particular include C_{10} - C_{18} alkyl dimethyl amine oxides and C_8 - C_{12} alkoxy ethyl dihydroxy ethyl amine oxides.

When included therein, the laundry detergent compositions of the present invention typically comprise from 0.2% to about 15%, preferably from about 1% to about 10% by weight of such semi-polar nonionic surfactants.

Builder system

30

The compositions according to the present invention may further comprise a builder system. Any conventional builder system is suitable for use herein including aluminosilicate materials, silicates, polycarboxylates and fatty acids, materials such as ethylenediamine tetraacetate, metal ion sequestrants such as aminopolyphosphonates, particularly ethylenediamine tetramethylene phosphonic acid and diethylene triamine pentamethylenephosphonic acid. Though less

preferred for obvious environmental reasons, phosphate builders can also be used herein.

Suitable builders can be an inorganic ion exchange material, commonly an inorganic hydrated aluminosilicate material, more particularly a hydrated synthetic zeolite such as hydrated zeolite A, X, B, HS or MAP.

Another suitable inorganic builder material is layered silicate, e.g. SKS-6 (Hoechst). SKS-6 is a crystalline layered silicate consisting of sodium silicate ($\text{Na}_2\text{Si}_2\text{O}_5$).

Suitable polycarboxylates containing one carboxy group include lactic acid, glycolic acid and ether derivatives thereof as disclosed in Belgian Patent Nos. 831,368, 821,369 and 821,370. Polycarboxylates containing two carboxy groups include the water-soluble salts of succinic acid, malonic acid, (ethylenedioxy) diacetic acid, maleic acid, diglycollic acid, tartaric acid, tartronic acid and fumaric acid, as well as the ether carboxylates described in German Offenle-enschrift 2,446,686, and 2,446,487, US 3,935,257 and the sulfinyl carboxylates described in Belgian Patent No. 840,623. Polycarboxylates containing three carboxy groups include, in particular, water-soluble citrates, aconitrates and citraconates as well as succinate derivatives such as the carboxymethyloxysuccinates described in British Patent No. 1,379,241, lactoxysuccinates described in Netherlands Application 7205873, and the oxypolycarboxylate materials such as 2-oxa-1,1,3-propane tricarboxylates described in British Patent No. 1,387,447.

Polycarboxylates containing four carboxy groups include oxydisuccinates disclosed in British Patent No. 1,261,829, 1,1,2,2,-ethane tetracarboxylates, 1,1,3,3-propane tetracarboxylates containing sulfo substituents include the sulfosuccinate derivatives disclosed in British Patent Nos. 1,398,421 and 1,398,422 and in US 3,936,448, and the sulfonated pyrolysed citrates described in British Patent

No. 1,082,179, while polycarboxylates containing phosphone substituents are disclosed in British Patent No. 1,439,000.

Alicyclic and heterocyclic polycarboxylates include
5 cyclopentane-cis,cis-cis-tetracarboxylates,
cyclopentadienide pentacarboxylates, 2,3,4,5-tetrahydro-
furan - cis, cis, cis-tetracarboxylates, 2,5-tetrahydro-
furan-cis, discarboxylates, 2,2,5,5,-tetrahydrofuran -
tetracarboxylates, 1,2,3,4,5,6-hexane - hexacarboxylates
10 and carboxymethyl derivatives of polyhydric alcohols such
as sorbitol, mannitol and xylitol. Aromatic
polycarboxylates include mellitic acid, pyromellitic acid
and the phthalic acid derivatives disclosed in British
Patent No. 1,425,343.

15

Of the above, the preferred polycarboxylates are hydroxy-
carboxylates containing up to three carboxy groups per
molecule, more particularly citrates.

20 Preferred builder systems for use in the present
compositions include a mixture of a water-insoluble
aluminosilicate builder such as zeolite A or of a layered
silicate (SKS-6), and a water-soluble carboxylate chelating
agent such as citric acid.

25

A suitable chelant for inclusion in the detergent composi-
tions in accordance with the invention is ethylenediamine-
N,N'-disuccinic acid (EDDS) or the alkali metal, alkaline
earth metal, ammonium, or substituted ammonium salts
30 thereof, or mixtures thereof. Preferred EDDS compounds are
the free acid form and the sodium or magnesium salt
thereof. Examples of such preferred sodium salts of EDDS
include Na_2EDDS and Na_4EDDS . Examples of such preferred
magnesium salts of EDDS include MgEDDS and Mg_2EDDS . The
35 magnesium salts are the most preferred for inclusion in
compositions in accordance with the invention.

Preferred builder systems include a mixture of a water-insoluble aluminosilicate builder such as zeolite A, and a water soluble carboxylate chelating agent such as citric acid.

5

Other builder materials that can form part of the builder system for use in granular compositions include inorganic materials such as alkali metal carbonates, bicarbonates, silicates, and organic materials such as the organic
10 phosphonates, amino polyalkylene phosphonates and amino polycarboxylates.

Other suitable water-soluble organic salts are the homo- or co-polymeric acids or their salts, in which the
15 polycarboxylic acid comprises at least two carboxyl radicals separated from each other by not more than two carbon atoms.

Polymers of this type are disclosed in GB-A-1,596,756.

20 Examples of such salts are polyacrylates of MW 2000-5000 and their copolymers with maleic anhydride, such copolymers having a molecular weight of from 20,000 to 70,000, especially about 40,000.

25 Detergency builder salts are normally included in amounts of from 5% to 80% by weight of the composition. Preferred levels of builder for liquid detergents are from 5% to 30%.

Enzymes

30

Preferred detergent compositions, in addition to the enzyme preparation of the invention, comprise other enzyme(s) which provides cleaning performance and/or fabric care benefits.

35

Such enzymes include proteases, lipases, cutinases, amylases, cellulases, peroxidases, oxidases (e.g. laccases).

Proteases: Any protease suitable for use in alkaline solutions can be used. Suitable proteases include those of animal, vegetable or microbial origin. Microbial origin is preferred. Chemically or genetically modified mutants are included. The protease may be a serine protease, preferably an alkaline microbial protease or a trypsin-like protease. Examples of alkaline proteases are subtilisins, especially those derived from Bacillus, e.g., subtilisin Novo, subtilisin Carlsberg, subtilisin 309, subtilisin 147 and subtilisin 168 (described in WO 89/06279). Examples of trypsin-like proteases are trypsin (e.g. of porcine or bovine origin) and the Fusarium protease described in WO 89/06270.

Preferred commercially available protease enzymes include those sold under the trade names Alcalase, Savinase, Primase, Durazym, and Esperase by Novo Nordisk A/S (Denmark), those sold under the tradename Maxatase, Maxacal, Maxapem, Properase, Purafect and Purafect OXP by Genencor International, and those sold under the tradename Opticlean and Optimase by Solvay Enzymes. Protease enzymes may be incorporated into the compositions in accordance with the invention at a level of from 0.00001% to 2% of enzyme protein by weight of the composition, preferably at a level of from 0.0001% to 1% of enzyme protein by weight of the composition, more preferably at a level of from 0.001% to 0.5% of enzyme protein by weight of the composition, even more preferably at a level of from 0.01% to 0.2% of enzyme protein by weight of the composition.

Lipases: Any lipase suitable for use in alkaline solutions can be used. Suitable lipases include those of bacterial or fungal origin. Chemically or genetically modified mutants are included.

Examples of useful lipases include a Humicola lanuginosa lipase, e.g., as described in EP 258 068 and EP 305 216, a

Rhizomucor miehei lipase, e.g., as described in EP 238 023, a Candida lipase, such as a C. antarctica lipase, e.g., the C. antarctica lipase A or B described in EP 214 761, a Pseudomonas lipase such as a P. alcaligenes and P. pseudoalcaligenes lipase, e.g., as described in EP 218 272, a P. cepacia lipase, e.g., as described in EP 331 376, a P. stutzeri lipase, e.g., as disclosed in GB 1,372,034, a P. fluorescens lipase, a Bacillus lipase, e.g., a B. subtilis lipase (Dartois et al., (1993), Biochemica et Biophysica acta 1131, 253-260), a B. stearothermophilus lipase (JP 64/744992) and a B. pumilus lipase (WO 91/16422).

Furthermore, a number of cloned lipases may be useful, including the Penicillium camembertii lipase described by Yamaguchi et al., (1991), Gene 103, 61-67), the Geotricum candidum lipase (Schimada, Y. et al., (1989), J. Biochem., 106, 383-388), and various Rhizopus lipases such as a R. delemar lipase (Hass, M.J et al., (1991), Gene 109, 117-113), a R. niveus lipase (Kugimiya et al., (1992), Biosci. Biotech. Biochem. 56, 716-719) and a R. oryzae lipase.

Other types of lipolytic enzymes such as cutinases may also be useful, e.g., a cutinase derived from Pseudomonas mendocina as described in WO 88/09367, or a cutinase derived from Fusarium solani pisi (e.g. described in WO 90/09446).

Especially suitable lipases are lipases such as M1 Lipase™, Luma fast™ and Lipomax™ (Genencor), Lipolase™ and Lipolase Ultra™ (Novo Nordisk A/S), and Lipase P "Amano" (Amano Pharmaceutical Co. Ltd.).

The lipases are normally incorporated in the detergent composition at a level of from 0.00001% to 2% of enzyme protein by weight of the composition, preferably at a level of from 0.0001% to 1% of enzyme protein by weight of the composition, more preferably at a level of from 0.001% to 0.5% of enzyme protein by weight of the composition, even more preferably at a level of from 0.01% to 0.2% of enzyme

protein by weight of the composition.

Amylases: Any amylase (α and/or β) suitable for use in alkaline solutions can be used. Suitable amylases include those of bacterial or fungal origin. Chemically or genetically modified mutants are included. Amylases include, for example, α -amylases obtained from a special strain of B. licheniformis, described in more detail in GB 1,296,839. Commercially available amylases are Duramyl™, Termamyl™, Fungamyl™ and BAN™ (available from Novo Nordisk A/S) and Rapidase™ and Maxamyl P™ (available from Genencor).

The amylases are normally incorporated in the detergent composition at a level of from 0.00001% to 2% of enzyme protein by weight of the composition, preferably at a level of from 0.0001% to 1% of enzyme protein by weight of the composition, more preferably at a level of from 0.001% to 0.5% of enzyme protein by weight of the composition, even more preferably at a level of from 0.01% to 0.2% of enzyme protein by weight of the composition.

Cellulases: Any cellulase suitable for use in alkaline solutions can be used. Suitable cellulases include those of bacterial or fungal origin. Chemically or genetically modified mutants are included. Suitable cellulases are disclosed in US 4,435,307, which discloses fungal cellulases produced from Humicola insolens. Especially suitable cellulases are the cellulases having colour care benefits. Examples of such cellulases are cellulases described in European patent application No. 0 495 257.

Commercially available cellulases include Celluzyme™ produced by a strain of Humicola insolens, (Novo Nordisk A/S), and KAC-500(B)™ (Kao Corporation).

Cellulases are normally incorporated in the detergent

composition at a level of from 0.00001% to 2% of enzyme protein by weight of the composition, preferably at a level of from 0.0001% to 1% of enzyme protein by weight of the composition, more preferably at a level of from 0.001% to 5 0.5% of enzyme protein by weight of the composition, even more preferably at a level of from 0.01% to 0.2% of enzyme protein by weight of the composition.

Peroxidases/Oxidases: Peroxidase enzymes are used in 10 combination with hydrogen peroxide or a source thereof (e.g. a percarbonate, perborate or persulfate). Oxidase enzymes are used in combination with oxygen. Both types of enzymes are used for "solution bleaching", i.e. to prevent transfer of a textile dye from a dyed fabric to another 15 fabric when said fabrics are washed together in a wash liquor, preferably together with an enhancing agent as described in e.g. WO 94/12621 and WO 95/01426. Suitable peroxidases/oxidases include those of plant, bacterial or fungal origin. Chemically or genetically modified mutants 20 are included.

Peroxidase and/or oxidase enzymes are normally incorporated in the detergent composition at a level of from 0.00001% to 2% of enzyme protein by weight of the composition, 25 preferably at a level of from 0.0001% to 1% of enzyme protein by weight of the composition, more preferably at a level of from 0.001% to 0.5% of enzyme protein by weight of the composition, even more preferably at a level of from 0.01% to 0.2% of enzyme protein by weight of the 30 composition.

Mixtures of the above mentioned enzymes are encompassed herein, in particular a mixture of a protease, an amylase, a lipase and/or a cellulase.

35

The enzyme of the invention, or any other enzyme incorporated in the detergent composition, is normally incorporated in the detergent composition at a level from

0.00001% to 2% of enzyme protein by weight of the composition, preferably at a level from 0.0001% to 1% of enzyme protein by weight of the composition, more preferably at a level from 0.001% to 0.5% of enzyme protein by weight of the composition, even more preferably at a level from 0.01% to 0.2% of enzyme protein by weight of the composition.

Bleaching agents: Additional optional detergent ingredients that can be included in the detergent compositions of the present invention include bleaching agents such as PB1, PB4 and percarbonate with a particle size of 400-800 microns. These bleaching agent components can include one or more oxygen bleaching agents and, depending upon the bleaching agent chosen, one or more bleach activators. When present oxygen bleaching compounds will typically be present at levels of from about 1% to about 25%. In general, bleaching compounds are optional added components in non-liquid formulations, e.g. granular detergents.

The bleaching agent component for use herein can be any of the bleaching agents useful for detergent compositions including oxygen bleaches as well as others known in the art.

The bleaching agent suitable for the present invention can be an activated or non-activated bleaching agent.

One category of oxygen bleaching agent that can be used encompasses percarboxylic acid bleaching agents and salts thereof. Suitable examples of this class of agents include magnesium monoperoxyphthalate hexahydrate, the magnesium salt of meta-chloro perbenzoic acid, 4-nonylamino-4-oxoperoxybutyric acid and diperoxydodecanedioic acid. Such bleaching agents are disclosed in US 4,483,781, US 740,446, EP 0 133 354 and US 4,412,934. Highly preferred bleaching agents also include 6-nonylamino-6-oxoperoxyacaproic acid as described in US 4,634,551.

Another category of bleaching agents that can be used encompasses the halogen bleaching agents. Examples of hypohalite bleaching agents, for example, include trichloro isocyanuric acid and the sodium and potassium dichloroisocyanurates and N-chloro and N-bromo alkane sulphonamides. Such materials are normally added at 0.5-10% by weight of the finished product, preferably 1-5% by weight.

10

The hydrogen peroxide releasing agents can be used in combination with bleach activators such as tetraacetythylenediamine (TAED), nonanoyloxybenzenesulfonate (NOBS, described in US 4,412,934), 3,5-trimethylhexanoyloxybenzenesulfonate (ISONOBS, described in EP 120 591) or pentaacetylglucose (PAG), which are perhydrolyzed to form a peracid as the active bleaching species, leading to improved bleaching effect. In addition, very suitable are the bleach activators C8(6-octanamido-caproyl) oxybenzene-sulfonate, C9(6-nonanamido caproyl) oxybenzenesulfonate and C10 (6-decanamido caproyl) oxybenzenesulfonate or mixtures thereof. Also suitable activators are acylated citrate esters such as disclosed in European Patent Application No. 91870207.7.

25

Useful bleaching agents, including peroxyacids and bleaching systems comprising bleach activators and peroxygen bleaching compounds for use in cleaning compositions according to the invention are described in application USSN 08/136,626.

30

The hydrogen peroxide may also be present by adding an enzymatic system (i.e. an enzyme and a substrate therefore) which is capable of generation of hydrogen peroxide at the beginning or during the washing and/or rinsing process. Such enzymatic systems are disclosed in European Patent Application EP 0 537 381.

35

Bleaching agents other than oxygen bleaching agents are also known in the art and can be utilized herein. One type of non-oxygen bleaching agent of particular interest
5 includes photoactivated bleaching agents such as the sulfonated zinc and/or aluminium phthalocyanines. These materials can be deposited upon the substrate during the washing process. Upon irradiation with light, in the presence of oxygen, such as by hanging clothes out to dry
10 in the daylight, the sulfonated zinc phthalocyanine is activated and, consequently, the substrate is bleached. Preferred zinc phthalocyanine and a photoactivated bleaching process are described in US 4,033,718. Typically, detergent composition will contain about 0.025% to about
15 1.25%, by weight, of sulfonated zinc phthalocyanine.

Bleaching agents may also comprise a manganese catalyst. The manganese catalyst may, e.g., be one of the compounds described in "Efficient manganese catalysts for low-
20 temperature bleaching", Nature 369, 1994, pp. 637-639.

Suds suppressors: Another optional ingredient is a suds suppressor, exemplified by silicones, and silica-silicone mixtures. Silicones can generally be represented by
25 alkylated polysiloxane materials, while silica is normally used in finely divided forms exemplified by silica aerogels and xerogels and hydrophobic silicas of various types. These materials can be incorporated as particulates, in which the suds suppressor is advantageously releasably
30 incorporated in a water-soluble or waterdispersible, substantially non surface-active detergent impermeable carrier. Alternatively the suds suppressor can be dissolved or dispersed in a liquid carrier and applied by spraying on to one or more of the other components.

35

A preferred silicone suds controlling agent is disclosed in US 3,933,672. Other particularly useful suds suppressors are the self-emulsifying silicone suds suppressors,

described in German Patent Application DTOS 2,646,126. An example of such a compound is DC-544, commercially available from Dow Corning, which is a siloxane-glycol copolymer. Especially preferred suds controlling agent are the suds suppressor system comprising a mixture of silicone oils and 2-alkyl-alkanols. Suitable 2-alkyl-alkanols are 2-butyl-octanol which are commercially available under the trade name Isofol 12 R.

10 Such suds suppressor system are described in European Patent Application EP 0 593 841.

Especially preferred silicone suds controlling agents are described in European Patent Application No. 92201649.8.

15 Said compositions can comprise a silicone/ silica mixture in combination with fumed nonporous silica such as Aerosil^R.

The suds suppressors described above are normally employed at levels of from 0.001% to 2% by weight of the composition, preferably from 0.01% to 1% by weight.

Other components: Other components used in detergent compositions may be employed such as soil-suspending agents, soil-releasing agents, optical brighteners, abrasives, bactericides, tarnish inhibitors, coloring agents, and/or encapsulated or nonencapsulated perfumes.

Especially suitable encapsulating materials are water soluble capsules which consist of a matrix of polysaccharide and polyhydroxy compounds such as described in GB 1,464,616.

Other suitable water soluble encapsulating materials comprise dextrans derived from ungelatinized starch acid esters of substituted dicarboxylic acids such as described in US 3,455,838. These acid-ester dextrans are, preferably, prepared from such starches as waxy maize, waxy sorghum,


sago, tapioca and potato. Suitable examples of said encapsulation materials include N-Lok manufactured by National Starch. The N-Lok encapsulating material consists of a modified maize starch and glucose. The starch is
5 modified by adding monofunctional substituted groups such as octenyl succinic acid anhydride.

Antiredeposition and soil suspension agents suitable herein include cellulose derivatives such as methylcellulose,
10 carboxymethylcellulose and hydroxyethylcellulose, and homo- or co-polymeric polycarboxylic acids or their salts. Polymers of this type include the polyacrylates and maleic anhydride-acrylic acid copolymers previously mentioned as builders, as well as copolymers of maleic anhydride with
15 ethylene, methylvinyl ether or methacrylic acid, the maleic anhydride constituting at least 20 mole percent of the copolymer. These materials are normally used at levels of from 0.5% to 10% by weight, more preferably from 0.75% to 8%, most preferably from 1% to 6% by weight of the
20 composition.

Preferred optical brighteners are anionic in character, examples of which are disodium 4,4'-bis-(2-diethanolamino-4-anilino -s- triazin-6-ylamino)stilbene-2:2' disulphonate,
25 disodium 4, - 4'-bis-(2-morpholino-4-anilino-s-triazin-6-ylamino-stilbene-2:2' - disulphonate, disodium 4,4' - bis-(2,4-dianilino-s-triazin-6-ylamino)stilbene-2:2' - disulphonate, monosodium 4',4'' - bis-(2,4-dianilino-s-triazin-6 ylamino)stilbene-2-sulphonate, disodium 4,4' -bis-
30 (2-anilino-4-(N-methyl-N-2-hydroxyethylamino)-s-triazin-6-ylamino)stilbene-2,2' - disulphonate, di-sodium 4,4' -bis-(4-phenyl-2,1,3-triazol-2-yl)-stilbene-2,2' disulphonate, di-so-dium 4,4'bis(2-anilino-4-(1-methyl-2-hydroxyethylamino)-s-triazin-6-ylami-no)stilbene-
35 2,2'disulphonate, sodium 2(stilbyl-4''-(naphtho-1',2':4,5)-1,2,3, - triazole-2''-sulphonate and 4,4'-bis(2-sulphostyryl)biphenyl.

Other useful polymeric materials are the polyethylene glycols, particularly those of molecular weight 1000-10000, more particularly 2000 to 8000 and most preferably about 4000. These are used at levels of from 0.20% to 5% more preferably from 0.25% to 2.5% by weight. These polymers and the previously mentioned homo- or co-polymeric polycarboxylate salts are valuable for improving whiteness maintenance, fabric ash deposition, and cleaning performance on clay, proteinaceous and oxidizable soils in the presence of transition metal impurities.

Soil release agents useful in compositions of the present invention are conventionally copolymers or terpolymers of terephthalic acid with ethylene glycol and/or propylene glycol units in various arrangements. Examples of such polymers are disclosed in US 4,116,885 and 4,711,730 and EP 0 272 033. A particular preferred polymer in accordance with EP 0 272 033 has the formula:



where PEG is $-(\text{OC}_2\text{H}_4)_n-$, PO is $(\text{OC}_3\text{H}_6\text{O})$ and T is $(\text{pOOC}_6\text{H}_4\text{CO})$.

Also very useful are modified polyesters as random copolymers of dimethyl terephthalate, dimethyl sulfoisophthalate, ethylene glycol and 1,2-propanediol, the end groups consisting primarily of sulphobenzoate and secondarily of mono esters of ethylene glycol and/or 1,2-propanediol. The target is to obtain a polymer capped at both end by sulphobenzoate groups, "primarily", in the present context most of said copolymers herein will be endcapped by sulphobenzoate groups. However, some copolymers will be less than fully capped, and therefore their end groups may consist of monoester of ethylene glycol and/or 1,2-propanediol, thereof consist "secondarily" of such species.

The selected polyesters herein contain about 46% by weight of dimethyl terephthalic acid, about 16% by weight of 1,2-propanediol, about 10% by weight ethylene glycol, about 13% by weight of dimethyl sulfobenzoic acid and about 15% by weight of sulfoisophthalic acid, and have a molecular weight of about 3.000. The polyesters and their method of preparation are described in detail in EP 311 342.

10

Softening agents: Fabric softening agents can also be incorporated into laundry detergent compositions in accordance with the present invention. These agents may be inorganic or organic in type. Inorganic softening agents are exemplified by the smectite clays disclosed in GB-A-1 400898 and in US 5,019,292. Organic fabric softening agents include the water insoluble tertiary amines as disclosed in GB-A1 514 276 and EP 0 011 340 and their combination with mono C₁₂-C₁₄ quaternary ammonium salts are disclosed in EP- B-0 026 528 and di-long-chain amides as disclosed in EP 0 242 919. Other useful organic ingredients of fabric softening systems include high molecular weight polyethylene oxide materials as disclosed in EP 0 299 575 and 0 313 146.

25

Levels of smectite clay are normally in the range from 5% to 15%, more preferably from 8% to 12% by weight, with the material being added as a dry mixed component to the remainder of the formulation. Organic fabric softening agents such as the water-insoluble tertiary amines or dilong chain amide materials are incorporated at levels of from 0.5% to 5% by weight, normally from 1% to 3% by weight whilst the high molecular weight polyethylene oxide materials and the water soluble cationic materials are added at levels of from 0.1% to 2%, normally from 0.15% to 1.5% by weight. These materials are normally added to the spray dried portion of the composition, although in some instances it may be more convenient to add them as a dry

mixed particulate, or spray them as molten liquid on to other solid components of the composition.

5 Polymeric dye-transfer inhibiting agents: The detergent compositions according to the present invention may also comprise from 0.001% to 10%, preferably from 0.01% to 2%, more preferably from 0.05% to 1% by weight of polymeric dye-transfer inhibiting agents. Said polymeric dye-
10 transfer inhibiting agents are normally incorporated into detergent compositions in order to inhibit the transfer of dyes from colored fabrics onto fabrics washed therewith. These polymers have the ability of complexing or adsorbing the fugitive dyes washed out of dyed fabrics before the
15 dyes have the opportunity to become attached to other articles in the wash.

Especially suitable polymeric dye-transfer inhibiting agents are polyamine N-oxide polymers, copolymers of N-
20 vinyl-pyrrolidone and N-vinylimidazole, polyvinyl-pyrrolidone polymers, polyvinylloxazolidones and polyvinylimidazoles or mixtures thereof.

Addition of such polymers also enhances the performance of
25 the enzymes according to the invention.

The detergent composition according to the invention can be in liquid, paste, gels, bars or granular forms.

Non-dusting granulates may be produced, e.g., as disclosed
30 in US 4,106,991 and 4,661,452 (both to Novo Industri A/S) and may optionally be coated by methods known in the art. Examples of waxy coating materials are poly(ethylene oxide) products (polyethyleneglycol, PEG) with mean molecular weights of 1000 to 20000; ethoxylated nonylphenols having
35 from 16 to 50 ethylene oxide units; ethoxylated fatty alcohols in which the alcohol contains from 12 to 20 carbon atoms and in which there are 15 to 80 ethylene oxide units; fatty alcohols; fatty acids; and mono- and di- and

triglycerides of fatty acids. Examples of film-forming coating materials suitable for application by fluid bed techniques are given in GB 1483591.

5 Granular compositions according to the present invention can also be in "compact form", i.e. they may have a relatively higher density than conventional granular detergents, i.e. from 550 to 950 g/l; in such case, the granular detergent compositions according to the present
10 invention will contain a lower amount of "Inorganic filler salt", compared to conventional granular detergents; typical filler salts are alkaline earth metal salts of sulphates and chlorides, typically sodium sulphate; "Compact" detergent typically comprise not more than 10%
15 filler salt. The liquid compositions according to the present invention can also be in "concentrated form", in such case, the liquid detergent compositions according to the present invention will contain a lower amount of water, compared to conventional liquid detergents. Typically, the
20 water content of the concentrated liquid detergent is less than 30%, more preferably less than 20%, most preferably less than 10% by weight of the detergent compositions.

The compositions of the invention may for example, be
25 formulated as hand and machine laundry detergent compositions including laundry additive compositions and compositions suitable for use in the pretreatment of stained fabrics, rinse added fabric softener compositions, and compositions for use in general household hard surface
30 cleaning operations and dishwashing operations.

The following examples are meant to exemplify compositions for the present invention, but are not necessarily meant to limit or otherwise define the scope of the invention.
35 In the detergent compositions, the abbreviated component identifications have the following meanings:

- LAS: Sodium linear C_{12} alkyl benzene sulphonate
- TAS: Sodium tallow alkyl sulphate
- 5 XYAS: Sodium $C_{1X} - C_{1Y}$ alkyl sulfate
- SS: Secondary soap surfactant of formula 2-butyl octanoic acid
- 10 25EY: A $C_{12} - C_{15}$ predominantly linear primary alcohol condensed with an average of Y moles of ethylene oxide
- 45EY: A $C_{14} - C_{15}$ predominantly linear primary alcohol
15 condensed with an average of Y moles of ethylene oxide
- XYEzs: $C_{1X} - C_{1Y}$ sodium alkyl sulfate condensed with an average of Z moles of ethylene oxide per mole
- 20 Nonionic: $C_{13} - C_{15}$ mixed ethoxylated/propoxylated fatty alcohol with an average degree of ethoxylation of 3.8 and an average degree of propoxylation of 4.5 sold under the tradename Plurafax LF404 by BASF GmbH
- 25 CFAA: $C_{12} - C_{14}$ alkyl N-methyl glucamide
- TFAA: $C_{16} - C_{18}$ alkyl N-methyl glucamide
- Silicate: Amorphous Sodium Silicate ($SiO_2:Na_2O$ ratio =
30 2.0)
- NaSKS-6: Crystalline layered silicate of formula $\delta-Na_2Si_2O_5$
- 35 Carbonate: Anhydrous sodium carbonate
- Phosphate: Sodium tripolyphosphate

MA/AA: Copolymer of 1:4 maleic/acrylic acid, average molecular weight about 80,000

5 Polyacrylate: Polyacrylate homopolymer with an average molecular weight of 8,000 sold under the tradename PA30 by BASF GmbH

Zeolite A: Hydrated Sodium Aluminosilicate of formula
10 $\text{Na}_{12}(\text{AlO}_2\text{SiO}_2)_{12} \cdot 27\text{H}_2\text{O}$ having a primary particle size in the range from 1 to 10 micrometers

Citrate: Tri-sodium citrate dihydrate

15 Citric: Citric Acid

Perborate: Anhydrous sodium perborate monohydrate bleach, empirical formula $\text{NaBO}_2 \cdot \text{H}_2\text{O}_2$

20 PB4: Anhydrous sodium perborate tetrahydrate

Percarbonate: Anhydrous sodium percarbonate bleach of empirical formula $2\text{Na}_2\text{CO}_3 \cdot 3\text{H}_2\text{O}_2$

25 TAED: Tetraacetyl ethylene diamine

CMC: Sodium carboxymethyl cellulose

DETPMP: Diethylene triamine penta (methylene phosphonic
30 acid), marketed by Monsanto under the Tradename Dequest 2060

PVP: Polyvinylpyrrolidone polymer

35 EDDS: Ethylenediamine-N, N'-disuccinic acid, [S,S] isomer in the form of the sodium salt

Suds Suppressor: 25% paraffin wax Mpt 50°C, 17%
hydrophobic silica, 58% paraffin oil

Granular Suds suppressor: 12% Silicone/silica, 18%
5 stearyl alcohol, 70% starch in granular form

Sulphate: Anhydrous sodium sulphate

HMWPEO: High molecular weight polyethylene oxide
10

TAE 25: Tallow alcohol ethoxylate (25)

Detergent Example I

15

A granular fabric cleaning composition in accordance with
the invention may be prepared as follows:

20	Sodium linear C ₁₂ alkyl benzene sulfonate	6.5
	Sodium sulfate	15.0
25	Zeolite A	26.0
	Sodium nitrilotriacetate	5.0
	Enzyme of the invention	0.1
30	PVP	0.5
	TAED	3.0
35	Boric acid	4.0
	Perborate	18.0
	Phenol sulphonate	0.1
40	Minors	Up to 100

Detergent Example II

A compact granular fabric cleaning composition (density 800 g/l) in accord with the invention may be prepared as follows:

5	45AS	8.0
	25E3S	2.0
	25E5	3.0
10	25E3	3.0
	TFAA	2.5
15	Zeolite A	17.0
	NaSKS-6	12.0
	Citric acid	3.0
20	Carbonate	7.0
	MA/AA	5.0
25	CMC	0.4
	Enzyme of the invention	0.1
	TAED	6.0
30	Percarbonate	22.0
	EDDS	0.3
35	Granular suds suppressor	3.5
	water/minors	Up to 100%

Detergent Example III

40

Granular fabric cleaning compositions in accordance with the invention which are especially useful in the laundering of coloured fabrics were prepared as follows:

	LAS	10.7	-
45	TAS	2.4	-
	TFAA	-	4.0
	45AS	3.1	10.0
	45E7	4.0	-
	25E3S	-	3.0

68E11	1.8	-
25E5	-	8.0
Citrate	15.0	7.0
Carbonate	-	10
5 Citric acid	2.5	3.0
Zeolite A	32.1	25.0
Na-SKS-6	-	9.0
MA/AA	5.0	5.0
DETPMP	0.2	0.8
10 Enzyme of the invention	0.10	0.05
Silicate	2.5	-
Sulphate	5.2	3.0
PVP	0.5	-
Poly (4-vinylpyridine)-N-	-	0.2
15 Oxide/copolymer of vinyl- imidazole and vinyl- pyrrolidone		
Perborate	1.0	-
Phenol sulfonate	0.2	-
20 Water/Minors	Up to 100%	

Detergent Example IV

Granular fabric cleaning compositions in accordance with
25 the invention which provide "Softening through the wash"
capability may be prepared as follows:

45AS	-	10.0
LAS	7.6	-
30 68AS	1.3	-
45E7	4.0	-
25E3	-	5.0
Coco-alkyl-dimethyl hydroxy- ethyl ammonium chloride	1.4	1.0
35 Citrate	5.0	3.0
Na-SKS-6	-	11.0
Zeolite A	15.0	15.0

MA/AA	4.0	4.0
DETPMP	0.4	0.4
Perborate	15.0	-
Percarbonate	-	15.0
5 TAED	5.0	5.0
Smectite clay	10.0	10.0
HMWPEO	-	0.1
Enzyme of the invention	0.10	0.05
Silicate	3.0	5.0
10 Carbonate	10.0	10.0
Granular suds suppressor	1.0	4.0
CMC	0.2	0.1
Water/Minors	Up to 100%	

15 Detergent Example V

Heavy duty liquid fabric cleaning compositions in accordance with the invention may be prepared as follows:

	I	II
20 LAS acid form	-	25.0
Citric acid	5.0	2.0
25AS acid form	8.0	-
25AE2S acid form	3.0	-
25AE7	8.0	-
25 CFAA	5	-
DETPMP	1.0	1.0
Fatty acid	8	-
Oleic acid	-	1.0
Ethanol	4.0	6.0
30 Propanediol	2.0	6.0
Enzyme of the invention	0.10	0.05
Coco-alkyl dimethyl hydroxy ethyl ammonium chloride	-	3.0
35 Smectite clay	-	5.0
PVP	2.0	-
Water / Minors	Up to 100%	

In a preferred embodiment, the detergent or cleaning composition comprises the enhancing agent in an amount effective for killing or inhibiting cells, preferably in an amount above 1 ppm, more preferably above 10 ppm.

When used for preservation of food, beverages, cosmetics such as lotions, creams, gels, ointments, soaps, shampoos, conditioners, antiperspirants, deodorants, mouth wash; contact lens products, enzyme formulations, or food ingredients, the composition used in the method of the present invention may be incorporated into the unpreserved food, beverages, cosmetics, contact lens products, food ingredients or antiinflammatory product in an amount effective for killing or inhibiting growing microbial cells.

Thus, the composition used in the method of the invention may be useful as a disinfectant, e.g. in the treatment of acne, infections in the eye or the mouth, skin infections; in antiperspirants or deodorants; in foot bath salts; for cleaning and disinfection of contact lenses, hard surfaces, teeth (oral care), wounds, bruises and the like.

The method of the invention may advantageously be carried out at a relatively high pH, since it is contemplated that the bacteriocidal activities are optimal at high pH values.

In general it is contemplated that the composition of the present invention is useful for cleaning, disinfecting or inhibiting microbial growth on any hard surface. Examples of surfaces, which may advantageously be contacted with the composition of the invention are surfaces of process equipment used e.g. in food processing plants, dairies, chemical or pharmaceutical process plants, water sanitation systems, paper pulp processing plants, water treatment plants, and cooling towers. The composition of the invention should be used in an amount, which is

effective for cleaning, disinfecting or inhibiting microbial growth on the surface in question.

Further, it is contemplated that the composition of the invention can advantageously be used in a cleaning-in-place (C.I.P.) system for cleaning of process equipment of any kind.

The invention is illustrated by the following non-limiting examples.

EXAMPLE 1

Antibacterial activity of *Coprinus cinereus* recombinant peroxidase using different enhancing agents

5

The antibacterial activity of *Coprinus cinereus*, IFO 8371, recombinant peroxidase (rCIP) available from Novo Nordisk A/S, DK-2880 Bagsvaerd, Denmark, has been tested in a phosphate buffer with the following enhancing agents:

10 sodium thiocyanate (NaSCN), potassium iodide (KI), 10-phenothiazine propionic acid (PPT), butyl syringate (BS) and 2,2' azinobis(3-ethylbenzothiazoline-6-sulfonate) (ABTS). The hydrogen peroxide was either generated by glucose oxidase or added directly in the concentration of 5
15 mM. In order to avoid interference with substrate components, the experiment was carried out in a buffer instead of in a growth substrate.

P. fluorescens (10^4 cfu/ml) was treated with rCIP (3
20 POXU/ml) and the different enhancing agents (5 mM) for 15 min (pH 6.0, 40°C) combined with glucoseoxidase/glucose or hydrogen peroxide, respectively. The bactericidal activity was determined by plate counting and by incubation in Malthus. The detection times measured by the Malthus
25 instrument were converted to cfu/ml by a standard curve.

Indirect Malthus measurements were used when enumerating total survival cells (Malthus Flexi M2060, Malthus Instrument Limited). 3 ml of growth medium was transferred
30 to the outer chamber of the indirect Malthus cells, and 0.5 ml of sterile KOH (0.1 M) was transferred to the inner chamber. The cell suspensions were after enzyme treatment transferred to the outer chamber of the Malthus cell. As cells are growing in the outer chamber they produce CO₂
35 which will dissolve in the KOH in the inner chamber and thereby change the conductance of the KOH. The amount of CO₂ formed by the respiring cells surviving the enzyme treatment was used for estimating the number of viable

cells. When the conductance change is measurable by the Malthus, a detection time (dt) will be recorded. The dt's were converted to colony counts by use of a calibration curve relating cfu/ml to dt (fig. 1).

5 The results are shown in fig. 1 as the number of cells which survive the treatment (average from both Malthus experiments and plate counts). Glucose oxidase and hydrogen peroxide in the used concentration have a bactericidal activity, decreasing the cell number from 10^4 cfu/ml to
10 approximately 10^2 - 10^3 cfu/ml, independent of enhancing agent.

rCIP had a statistical significant bactericidal activity against *P. fluorescens* when NaSCN or KI were used as
15 enhancing agents. PPT had a slight bactericidal activity, however, only statistically significant at a 10% level. When KI was used as enhancing agent the activity was 100% bactericidal, both when determined by plate counts and Malthus.

20

EXAMPLE 2

**Antibacterial activity of rCIP and KI against *P. aeruginosa*
25 and *S. aureus*.**

The antibacterial activity of rCIP was tested in phosphate buffer (pH 6.0) against *Pseudomonas aeruginosa* ATCC 10146 and *Staphylococcus aureus* ATCC 25923 with potassium iodide
30 as electron donor, and hydrogen peroxide was added as electron acceptor. The cells (approximately 10^7 cfu/ml) were incubated with enzyme for 15 min at 40°C, and the experiment was carried out using a 2^3 factorial design reproduced two times. The bactericidal activity was
35 determined by incubation in Malthus. The detection times measured by the Malthus instrument were converted to cfu/ml by a standard curve.

Results for *P. aeruginosa*:

rCIP (POXU/ml)	H ₂ O ₂ (mM)	KI (mM)	survivors (cfu/ml)
0	0	0	1.7×10^7
0	0	0.5	2.1×10^7
0	0.5	0	1.2×10^7
0	0.5	0.5	6.6×10^6
0.5	0	0	2.6×10^7
0.5	0	0.5	1.7×10^7
0.5	0.5	0	9.0×10^6
0.5	0.5	0.5	1.0×10^5

- 5 The combination of rCIP with KI and hydrogen peroxide killed 99.4% of *P. aeruginosa*, which is a significant effect of the peroxidase compared to the combination of hydrogen peroxide and KI without peroxidase.

10

Results for *S. aureus*:

rCIP (POXU/ml)	H ₂ O ₂ (mM)	KI (mM)	survivors (cfu/ml)
0	0	0	2.9×10^8
0	0	0.5	2.0×10^8
0	0.5	0	1.3×10^8
0	0.5	0.5	1.4×10^8
0.5	0	0	1.6×10^8
0.5	0	0.5	3.4×10^8
0.5	0.5	0	1.8×10^8
0.5	0.5	0.5	5.6×10^6

The bactericidal activity against *S. aureus* was signifi-

cant for the combination of peroxidase, hydrogen peroxide and potassium iodide, whereas no activity of hydrogen peroxide combined with iodide was observed. The activity was observed as a 98% reduction in cfu/ml.

5

Increasing the concentration of peroxidase to 1 POXU/ml and the concentration of hydrogen peroxide and potassium iodide to 1 mM caused a total kill of the cell suspension.

5 SEQUENCE LISTING

INFORMATION FOR SEQ ID NO:1:

(i) SEQUENCE CHARACTERISTICS:

- 10 (A) LENGTH: 1306 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

15 (ii) MOLECULE TYPE: cDNA

(vi) ORIGINAL SOURCE:

- (A) ORGANISM: *Coprinus cinereus*
(B) STRAIN: IFO 8371

20

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:1:

ACTATGAAGC TCTCGCTTTT GTCCACCTTC GCTGCTGTCA TCATCGGTGC CCTCGCTCTA 60
25 CCCCAGGGTC CTGGAGGAGG CGGGTCAGTC ACTTGCCCCG GTGGACAGTC CACTTCGAAC 120
AGCCAGTGCT GCGTCTGGTT CGACGTTCTA GACGATCTTC AGACCAACTT CTACCAAGGG 180
TCCAAGTGTG AGAGCCCTGT TCGCAAGATT CTTAGAATTG TTTTCCATGA CGCGATCGGA 240
30 TTTTCGCCGG CGTTGACTGC TGCTGGTCAA TTCGGTGGTG GAGGAGCTGA TGGCTCCATC 300
ATTGCGCATT CGAACATCGA ATTGGCCTTC CCGGCTAATG GCGGCCTCAC CGACACCGTC 360
35 GAAGCCCTCC GCGCGGTCCG TATCAACCAC GGTGTCTCTT TCGGCGATCT CATCCAATTC 420
GCCACTGCCG TCGGCATGTC CAACTGCCCT GGCTCTCCCC GACTTGAGTT CTTGACGGGC 480

5
AGGAGCAACA GTTCCCAACC CTCCCCTCCT TCGTTGATCC CCGGTCCCGG AAACACTGTC 540
ACTGCTATCT TGGATCGTAT GGGCGATGCA GGCTTCAGCC CTGATGAAGT AGTTGACTTG 600
10 CTTGCTGCGC ATAGTTTGGC TTCTCAGGAG GGTTCGAACT CGGCCATCTT CAGGTCTCCT 660
TTGGACTCGA CCCCTCAAGT TTTCGATACC CAGTTCTACA TTGAGACCTT GCTCAAGGGT 720
ACCACTCAGC CTGGCCCTTC TCTCGGCTTT GCAGAGGAGC TCTCCCCCTT CCCTGGCGAA 780
15 TTCCGCATGA GGTCCGATGC TCTCTTGGCT CGCGACTCCC GAACCGCCTG CCGATGGCAA 840
TCCATGACCA GCAGCAATGA AGTTATGGGC CAGCGATACC GCGCCGCCAT GGCCAAGATG 900
20 TCTGTTCTCG GCTTCGACAG GAACGCCCTC ACCGATTGCT CTGACGTTAT TCCTTCTGCT 960
GTGTCCAACA ACGCTGCTCC TGTATCCCT GGTGGCCTTA CTGTCCGATGA TATCGAGGTT 1020
TCGTGCCCCA GCGAGCCTTT CCCTGAAATT GCTACCGCCT CAGGCCCTCT CCCCTCCCTC 1080
25 GCTCCTGCTC CTTGATCTGG TGAAGATGGT ACATCCTGCT CTCTCATCAT CCCTCTTAGC 1140
TATTTATCCA ATCTATCTAC CTATCTATGC AGTTTCTGTT CTATCACCAC AGGAAGCAAG 1200
30 AAAGAAAAAC AACAATGCAA CGTGAGCAGA AATCAGCAAA AAAATAAATC AGTATACTAC 1260
AGTAATGAGG CCAGTTTGCG TGGTGTGAGA AGTAAGTACG ACTCGG 1306

5 CLAIMS

1. An enzymatic antimicrobial composition comprising a peroxidase produced by or derivable from the fungus *Coprinus*, an enhancing agent and hydrogen peroxide or a source of
10 hydrogen peroxide.

2. The composition according to claim 1, wherein the peroxidase is a recombinant enzyme obtainable from *Coprinus cinereus*.

15

3. The composition according to claim 1 or 2, wherein the peroxidase is obtainable from *Coprinus cinereus*, IFO 8371, or is immunologically cross-reactive with the peroxidase obtainable from *Coprinus cinereus*, IFO 8371.

20

4. The composition according to any of the claims 1-3, wherein the source of hydrogen peroxide is an enzymatic hydrogen peroxide-generating system.

25 5. The composition according to claim 4, wherein the enzymatic system is selected from the group consisting of glucose oxidase/glucose, hexose oxidase/hexose, L- or D-amino acid oxidase/L- or D-amino acid, and lactate oxidase/lactate.


30 6. The composition according to any of the claims 1-5, wherein the enhancing agent is an electron donor.

7. The composition according to claim 6, wherein the electron donor is a water-soluble halide or thiocyanate salt.

35

5 8. The composition according to claim 6, wherein the enhancing agent is a compound having the formula:

10




15 in which formula X represents (-O-) or (-S-), and the substituent groups R^1 - R^9 , which may be identical or different, independently represents any of the following radicals: hydrogen, halogen, hydroxy, formyl, carboxy, and esters and salts hereof, carbamoyl, sulfo, and esters and salts hereof,
 20 sulfamoyl, nitro, amino, phenyl, C_1 - C_{14} -alkyl, C_1 - C_5 -alkoxy, carbonyl- C_1 - C_5 -alkyl, aryl- C_1 - C_5 -alkyl; which carbamoyl, sulfamoyl, and amino groups may furthermore be unsubstituted or substituted once or twice with a substituent group R^{10} ; and which phenyl may furthermore be unsubstituted or substituted
 25 with one or more substituent groups R^{10} ; and which C_1 - C_{14} -alkyl, C_1 - C_5 -alkoxy, carbonyl- C_1 - C_5 -alkyl, and aryl- C_1 - C_5 -alkyl groups may be saturated or unsaturated, branched or unbranched, and may furthermore be unsubstituted or substituted with one or more substituent groups R^{10} ;
 30 which substituent group R^{10} represents any of the following radicals: halogen, hydroxy, formyl, carboxy and esters and salts hereof, carbamoyl, sulfo and esters and salts hereof, sulfamoyl, nitro, amino, phenyl, aminoalkyl, piperidino, piperazinyl, pyrrolidin-1-yl, C_1 - C_5 -alkyl, C_1 - C_5 -alkoxy; which
 35 carbamoyl, sulfamoyl, and amino groups may furthermore be

5 unsubstituted or substituted once or twice with hydroxy, C₁-C₅-alkyl, C₁-C₅-alkoxy; and which phenyl may furthermore be substituted with one or more of the following radicals: halogen, hydroxy, amino, formyl, carboxy and esters and salts hereof, carbamoyl, sulfo and esters and salts hereof, and
10 sulfamoyl; and which C₁-C₅-alkyl, and C₁-C₅-alkoxy groups may furthermore be saturated or unsaturated, branched or unbranched, and may furthermore be substituted once or twice with any of the following radicals: halogen, hydroxy, amino, formyl, carboxy and esters and salts hereof, carbamoyl, sulfo
15 and esters and salts hereof, and sulfamoyl;
or in which general formula two of the substituent groups R¹-R⁹ may together form a group -B-, in which B represents any of the following the groups: (-CHR¹⁰-N=N-), (-CH=CH-)_n, (-CH=N-)_n or (-N=CR¹⁰-NR¹¹-), in which groups n-represents an
20 integer of from 1 to 3, R¹⁰ is a substituent group as defined above and R¹¹ is defined as R¹⁰.

9. The composition according to claim 8, wherein the enhancing agent is selected from the group consisting of
25 10-methylphenothiazine, phenothiazine-10-propionic acid, N-hydroxysuccinimide phenothiazine-10-propionate, 10-ethylphenothiazine-4-carboxylic acid, 10-ethylphenothiazine, 10-propylphenothiazine, 10-isopropylphenothiazine, methyl phenothiazine-10-propionate, 10-phenylphenothiazine, 10-allyl-
30 phenothiazine, 10-(3-(4-methylpiperazin-1-yl)propyl)phenothiazine, 10-(2-pyrrolidin-1-yl-ethyl)phenothiazine, 2-methoxy-10-methyl-phenothiazine, 1-methoxy-10-methylphenothiazine, 3-methoxy-10-methylphenothiazine, 3,10-dimethylphenothiazine, 3,7,10-trimethylphenothiazine, 10-(2-
35 hydroxyethyl)phenothiazine, 10-(3-

5 hydroxypropyl)phenothiazine, 3-(2-hydroxyethyl)-10-methylphenothiazine, 3-hydroxymethyl-10-methylphenothiazine, 3,7-dibromophenothiazine-10-propionic acid, phenothiazine-10-propionamide, chlorpromazine, 2-chloro-10-methylphenothiazine, 2-acetyl-10-methylphenothiazine, 10-10 methylphenoxazine, 10-ethylphenoxazine, phenoxazine-10-propionic acid, 10-(2-hydroxyethyl)phenoxazine and 4-carboxyphenoxazine-10-propionic acid.


10. The composition according to claim 6, wherein the
15 enhancing agent is a compound having the formula:



25 wherein A denotes a group -D, -CH=CH-D, -CH=CH-CH=CH-D, -CH=N-D, -N=N-D, or -N=CH-D; D is selected from the group consisting of -CO-E, -SO₂-E, -N-XY, and -N⁺-XYZ; E is -H, -OH, -R, or -OR, and X and Y and Z may be identical or different and selected from -H and -R; R is C₁-C₁₆ alkyl, preferably
30 saturated or unsaturated, branched or unbranched C₁-C₈ alkyl, optionally substituted with a carboxy, sulfo or amino group; and B and C may be the same or different and selected from C_mH_{2m+1}; 1 ≤ m ≤ 5.

11. The composition according to claim 10, wherein the enhancing agent is acetosyringone, methylsyringate, ethylsyringate, propylsyringate, butylsyringate, hexylsyringate, or octylsyringate.


12. The composition according to claim 6, wherein the enhancing agent is an azino compound having the general formula



in which formula the symbols A and B, which may be identical or different, independently represent any of the substituents II, III, IV, and V,


20

(II)




25

(III)




30

(IV)



(V)



- 5 in which substituents the symbols X and Y, which may be identical or different, independently represent carbon, nitrogen, which nitrogen may be unsubstituted or substituted with a substituent group R⁵, sulfur, oxygen, selenium or tellurium;
- 10 and in which substituents the substituent groups R¹, R², R³, and R⁴, which may be identical or different, independently represent hydrogen, halogen, a hydroxy group, a C₁-C₃ alkoxy group, a formyl group, a carboxy group, a sulfo group, a nitro group, a C₁-C₅ alkyl group, which alkyl group may
- 15 furthermore be saturated or unsaturated, linear or branched, or an amino group, which amino group may furthermore be unsubstituted or substituted once or twice with a substituent group R⁵;
- which substituent group R⁵ represents halogen, a hydroxy
- 20 group, a C₁-C₃ alkoxy group, a C₁-C₅ alkyl group, or an amino group; either in free form or in the form of an addition salt.

13. The composition according to claim 12, wherein the

25 enhancing agent is 2,2'-azino-bis(3-ethylbenzothiazoline-6-sulfonate), abbreviated ABTS.

14. A method of inhibiting microorganisms present in laundry, wherein the laundry is treated with a soaking, washing or

30 rinsing liquor comprising an effective amount of the composition according to any of the claims 1-13.

15. The method according to claim 14, wherein the laundry is treated in a washing machine.

5 16. A method of killing microbial cells present on human or
animal skin, hair, oral cavity, mucous membranes, teeth,
wounds, bruises or in the eye or inhibiting the growth
thereof, wherein the cells to be killed or inhibited or the
skin, mucous membrane, teeth, wound or bruise are/is con-
10 tacted with a composition according to any of the claims 1-
13.

17. The method according to claim 16, wherein the composition
is a liquid composition having disinfectant or preserving
15 properties such as a mouth wash composition, an
antiinflammatory liquid (a disinfectant), an eye lotion, a
perspirant, a deodorant, and a nasal spray; or a solid
composition having disinfectant or preserving properties such
as an eye ointment, an antiinflammatory ointment or cream, a
20 foot bath salt, a perspirant, and a deodorant.

18. A method of preserving a cosmetic product, wherein an
effective amount of the composition according to any of the
claims 1-13 is incorporated into the cosmetic product.
25

19. The method according to claim 18, wherein the cosmetic
product is a mouth wash composition, a cosmetic liquid or gel
or paste, an eye lotion, a perspirant, a deodorant, a nasal
spray, an eye ointment, an ointment or cream, a foot bath
30 salt.

20. Use of the composition according to any of the claims 1-
13 for cleaning or disinfection of contact lenses.

5 21. A method of cleaning, disinfecting or inhibiting microbial growth on a hard surface, wherein the surface is contacted with the composition according to any of the claims 1-13.

22. The method according to claim 21, wherein the hard
10 surface is a process equipment member of a cooling tower, a water treatment plant, a dairy, a food processing plant, a chemical or pharmaceutical process plant.

23. The method according to claim 21, wherein the hard
15 surface is a surface of water sanitation equipment.

24. The method according to claim 21, wherein the hard surface is a surface of equipment for paper pulp processing.

20 25. Use of the composition according to any of the claims 1-13 in a cleaning-in-place (C-I-P) system.

1/1




Fig. 1

INTERNATIONAL SEARCH REPORT

International application No.

PCT/DK 97/00205

A. CLASSIFICATION OF SUBJECT MATTER

IPC6: A01N 63/00, A01N 63/04, A01N 59/00, A01N 43/84, A01N 37/10, C11D 3/386, C11D 3/48

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC6: A01N, C11D

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

SE,DK,FI,NO classes as above

Electronic data base consulted during the international search (name of data base and, where practicable, search terms used)

CA,WPI

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	WO 9404127 A1 (MONTGOMERY, ROBERT, E.), 3 March 1994 (03.03.94), page 8; page 9 --	1-7,14,15, 18-25
X	EP 0500387 A2 (EXOHEMIS, INC.), 26 August 1992 (26.08.92), page 3, line 56; page 4, line 48 - line 51, claims 14,23,34-37 --	1-7,14,15, 18-25
X	WO 9527046 A2 (UNILEVER N.V.), 12 October 1995 (12.10.95), claims 1-9 --	1-7,14,15, 18-25
X	US 4996146 A (JACK H. KESSLER), 26 February 1991 (26.02.91) --	1-7,14,15, 18-25



Further documents are listed in the continuation of Box C.



See patent family annex.

* Special categories of cited documents:

- "A" document defining the general state of the art which is not considered to be of particular relevance
- "E" earlier document but published on or after the international filing date
- "L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- "O" document referring to an oral disclosure, use, exhibition or other means
- "P" document published prior to the international filing date but later than the priority date claimed

"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

"X" document of particular relevance: the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

"Y" document of particular relevance: the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art

"&" document member of the same patent family

Date of the actual completion of the international search

4 July 1997

Date of mailing of the international search report

08-07- 1997

Name and mailing address of the ISA/

Swedish Patent Office

Box 5055, S-102 42 STOCKHOLM

Facsimile No. +46 8 666 02 86

Authorized officer

Gerd Strandell

Telephone No. +46 8 782 25 00

INTERNATIONAL SEARCH REPORT

International application No.

PCT/DK 97/00205

C (Continuation). DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	US 4588586 A (JACK H. KESSLER ET AL), 13 May 1986 (13.05.86) --	1-7,14,15, 18-25
X	US 5227161 A (JACK H. KESSLER), 13 July 1993 (13.07.93) --	1-7,14,15, 18-25
X	WO 9201466 A1 (UNIVERSITE LIBRE DE BRUXELLES), 6 February 1992 (06.02.92), claims 1,2,6-10,12-13, 16-18,21-23 --	1-7,14,15, 18-25
X	WO 9405252 A1 (MONTGOMERY, ROBERT, E.), 17 March 1994 (17.03.94), the claims --	1-7,14,15, 18-25
X	WO 9610079 A1 (NOVO NORDISK A/S), 4 April 1996 (04.04.96), page 6, line 1 - line 5; page 6, line 30 - line 33, claims 1-5,12-16,20-24 --	1-6,10,11
X	WO 9412621 A1 (NOVO NORDISK), 9 June 1994 (09.06.94), page 7, line 30 - page 8, line 29; page 20, line 23 - line 27; page 21, line 23 - line 27, figures 1A-1E, claims 1,6-8,23, 28-30,37,42-44 --	1-6,8,9
X	WO 9324618 A1 (NOVO NORDISK A/S), 9 December 1993 (09.12.93), page 5, line 5 - line 12, claims 11-23 --	1-6
X	WO 9510602 A1 (NOVO NORDISK A/S), 20 April 1995 (20.04.95), claims 5-12 -- -----	1-6,8,9

INTERNATIONAL SEARCH REPORT

International application No.

PCT/DK 97/00205

Box I Observations where certain claims were found unsearchable (Continuation of item 1 of first sheet)

This international search report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:

1. ☒ Claims Nos.: 16, 17
because they relate to subject matter not required to be searched by this Authority, namely:
See PCT Rule 39.1(iv): Methods for treatment of the human or animal body by surgery or therapy, as well as diagnostic methods.
2. ☐ Claims Nos.:
because they relate to parts of the international application that do not comply with the prescribed requirements to such an extent that no meaningful international search can be carried out, specifically:
3. ☐ Claims Nos.:
because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 6.4(a).

Box II Observations where unity of invention is lacking (Continuation of item 2 of first sheet)

This International Searching Authority found multiple inventions in this international application, as follows:

1. ☐ As all required additional search fees were timely paid by the applicant, this international search report covers all searchable claims.
2. ☐ As all searchable claims could be searched without effort justifying an additional fee, this Authority did not invite payment of any additional fee.
3. ☐ As only some of the required additional search fees were timely paid by the applicant, this international search report covers only those claims for which fees were paid, specifically claims Nos.:
4. ☐ No required additional search fees were timely paid by the applicant. Consequently, this international search report is restricted to the invention first mentioned in the claims; it is covered by claims Nos.:

Remark on Protest

☐

The additional search fees were accompanied by the applicant's protest.

☐

No protest accompanied the payment of additional search fees.

INTERNATIONAL SEARCH REPORT

Information on patent family members

03/06/97

International application No.

PCT/DK 97/00205

Patent document cited in search report			Publication date	Patent family member(s)		Publication date
WO	9404127	A1	03/03/94	AU	5089193 A	15/03/94
				EP	0746303 A	11/12/96
				US	5262151 A	16/11/93
EP	0500387	A2	26/08/92	AU	663869 B	26/10/95
				AU	1536492 A	15/09/92
				CA	2061601 A	22/08/92
				IL	100997 A	12/09/96
				JP	6505482 T	23/06/94
				US	5389369 A	14/02/95
				US	5451402 A	19/09/95
				US	5510104 A	23/04/96
				US	5565197 A	15/10/96
				WO	9214484 A	03/09/92
WO	9527046	A2	12/10/95	AU	2085995 A	23/10/95
				AU	2215495 A	23/10/95
				CA	2182966 A	12/10/95
				CN	1146782 A	02/04/97
				EP	0753055 A	15/01/97
				NL	9401048 A	01/11/95
				PL	316571 A	20/01/97
				WO	9527009 A	12/10/95
US	4996146	A	26/02/91	NONE		
US	4588586	A	13/05/86	US	4473550 A	25/09/84
US	5227161	A	13/07/93	US	5370815 A	06/12/94
WO	9201466	A1	06/02/92	AT	123950 T	15/07/95
				DE	69110678 D,T	14/03/96
				EP	0540547 A,B	12/05/93
				JP	6501453 T	17/02/94
				US	5503853 A	02/04/96
WO	9405252	A1	17/03/94	AU	4839893 A	29/03/94
				CA	2143111 A	17/03/94
				EP	0658096 A	21/06/95
				US	5310541 A	10/05/94
WO	9610079	A1	04/04/96	AU	3517695 A	19/04/96

INTERNATIONAL SEARCH REPORT

Information on patent family members

03/06/97

International application No.

PCT/DK 97/00205

Patent document cited in search report			Publication date	Patent family member(s)		Publication date
WO	9412621	A1	09/06/94	CA	2150562 A	09/06/94
				EP	0679183 A	02/11/95
				FI	952648 A	28/07/95
				JP	8506009 T	02/07/96
				AU	7937194 A	22/05/95
				CA	2175047 A	04/05/95
				EP	0730641 A	11/09/96
				WO	9511964 A	04/05/95
				CA	2150563 A	09/06/94
				EP	0677102 A	18/10/95
				FI	952647 A	31/05/95
				JP	8503371 T	16/04/96
				WO	9412620 A	09/06/94
WO	324618	A1	09/12/93	NONE		
WO	9510602	A1	20/04/95	AU	7853194 A	04/05/95
				CN	1133062 A	09/10/96
				EP	0724631 A	07/08/96
				FI	961650 A	15/04/96